Nefrologia xx (2025) 501391







journal homepage: www.revistanefrologia.com

Original article

Factors influencing the removal of protein-bound uremic toxins in hemodiafiltration

Factores que influyen en la depuración de toxinas urémicas unidas a proteínas en hemodiafiltración

Víctor Joaquín Escudero-Saiz o a,b, Elena Cuadrado-Payán Anría Rodríguez-García Gregori Casals Lida María Rodas Néstor Fontseré, José Jesús Broseta Broseta, Francisco Maduell o a,b,*

ARTICLE INFO

Keywords: Protein-bound uremic toxins Hemodiafiltration Convection P-cresyl sulfate Indoxyl sulfate

ABSTRACT

Introduction: Protein-bound uremic toxins (PBUTs) have a high affinity for albumin and they are associated with increased cardiovascular morbidity and mortality in hemodialysis patients. Among them, p-cresyl sulfate (pCS) and indoxyl sulfate (IS) stand out due to their high toxicity. Postdilution hemodiafiltration (HDF) is one of the dialysis techniques that has shown the greatest benefits in terms of patient survival.

Materials and methods: This observational, single-center, cross-sectional study evaluated PBUT clearance in 137 patients undergoing post-dilution HDF, analyzing the factors that influence their removal. Reduction ratios (RRs) of IS and pCS were measured, as well as their correlation with dialysis parameters and clinical variables

Results: The mean RR for IS was 53.4% \pm 9.3%, and for pCS, 48.2% \pm 11.3%. A significant correlation was observed between the RR of both toxins (r = 0.606; P < .01), suggesting similar elimination mechanisms. In addition, total convective volume showed a positive correlation with the RR of pCS (r = 0.19; P = .027) and a weak correlation with the RR of IS (r = 0.155; P = .07). A significant difference in clearance was found according to sex, with higher RRs in women (P < .001 for IS and P = .008 for pCS).

Conclusions: The clearance of PBUTs is primarily diffusive. Enhancing all variables related to this physical principle will improve the elimination of these toxins. Postdilution HDF with high convective volume slightly increases this clearance. However, the results remain insufficient given the high toxicity of these molecules. New strategies, such as the use of adsorptive membranes and competitive molecules, are needed to optimize their removal and reduce the negative cardiovascular impact in hemodialysis patients.

RESUMEN

Palabras clave:
Toxinas urémicas unidas a proteínas
Hemodiafiltración
Convección
Sulfato de indoxilo
Sulfato de p-cresol

Introducción: Las toxinas urémicas unidas a proteínas (PBUT) presentan una elevada afinidad por la albúmina y se relacionan con un incremento en la morbilidad y mortalidad cardiovascular en pacientes en hemodiálisis. Entre ellas, el p-cresil sulfato (pCS) y el indoxil sulfato (IS) destacan por su elevada toxicidad. La hemodiafiltración postdilucional (HDF) es una de la técnica de hemodiálisis que mayores beneficios ha demostrado en la supervivencia de los pacientes en hemodiálisis.

Material y métodos: Este estudio observacional, unicéntrico y transversal evaluó la depuración de PBUT en 137 pacientes en HDF postdilucional, analizando los factores que influyen en su eliminación. Se midieron los porcentajes de reducción (PR) de IS y pCS, así como su correlación con parámetros de diálisis y variables clínicas.

Resultados: Los resultados mostraron que el PR medio de IS fue del 53,4% \pm 9,3% y el de pCS del 48,2% \pm 11,3%. Se observó una correlación significativa entre el PR de ambas toxinas (r=0,606; P<,01), sugiriendo

https://doi.org/10.1016/j.nefroe.2025.501391

Received 28 April 2025; Accepted 18 July 2025

Available online xxx

2013-2514/© 2025 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

Please cite this article in press as: V.J. Escudero-Saiz, E. Cuadrado-Payán, M. Rodríguez-García, et al., Factors influencing the removal of protein-bound uremic toxins in hemodiafiltration, Nefrologia, https://doi.org/10.1016/j.nefroe.2025.501391

^a Servicio de Nefrología y Trasplante Renal, Hospital Clínic, Barcelona, Spain

^b Medicine Department, Faculty of Medicine and Health Sciences, Universitat de Barcelona, Barcelona, Spain

^c Servicio de Bioquímica y Genética Molecular, Hospital Clínic, Barcelona, Spain

^{*} Corresponding author.

E-mail address: fmaduell@clinic.cat (F. Maduell).

V.J. Escudero-Saiz, E. Cuadrado-Payán, M. Rodríguez-García et al.

Nefrologia xx (2025) 501391

mecanismos de eliminación similares. Además, el volumen convectivo total mostró una correlación positiva con el PR de pCS (r=0.19; P=.027) y una correlación débil con el PR de IS (r=0.155; P=.07). Se encontró una diferencia significativa en la depuración según el sexo, con mayores PR en mujeres (P<.001 para IS y P=.008 para pCS).

Conclusiones: La depuración de PBUT es fundamentalmente difusiva. Incrementar todas las variables relacionadas con este principio físico, aumentará la eliminación de dichas toxinas. La HDF postdilucional con alto volumen convectivo incrementa discretamente esta depuración. No obstante, los resultados siguen siendo insuficientes dada la alta toxicidad de estas moléculas. Se requieren nuevas estrategias, como el uso de membranas adsortivas y moléculas competidoras, para optimizar su depuración y reducir el impacto negativo en la salud cardiovascular de los pacientes en hemodiálisis.

Introduction

Uremic toxins are classified according to their physicochemical characteristics and purification capacity according to conventional hemodialysis techniques, following the current classification of the European Uremic Toxins Working Group (EUTox). Of these, proteinbound uremic toxins (PBUTs) include various groups of molecules^{2,3} with high affinity for plasma proteins, mainly albumin.⁴ These toxins are produced after the hepatic metabolism of precursors formed by the metabolism of dietary proteins of the intestinal microbiota; subsequently, PBUTs are eliminated by the kidneys by tubular secretion.⁵ PBUTs are associated with increased cardiovascular morbidity and mortality in hemodialysis patients.^{6–8} Among these compounds, pcresyl sulfate (pCS) and indoxyl sulfate (IS) are the most studied and have demonstrated the greatest toxicity to the cardiovascular system.³ In fact, pCS has the highest index of uremic toxicity (grade 4), which affects up to 7 organs, whereas has a toxicity grade of 3, with 6 organs affected.9,10

Significant variability in the percentage reduction (PR) of PBUT exists. 10 Conventional techniques of hemodialysis using high-flux hemodialysis, postdilution hemodiafiltration (HDF) 11 or expanded hemodialysis 12 present clearance rates lower than 55% and 50% for IS and pCS, respectively. 10 Owing to the small molecular weight of PBUT and its fundamentally diffusive depuration, extended-time techniques are the best available strategy for achieving the greatest degree of depuration. 13,14 Currently, other strategies are being developed, such as the use of competing molecules or the development of membranes with adsorptive capacity. 15

The objective of this study was to evaluate the clearance of PBUT in our dialysis unit, in postdilution HDF mode, and to evaluate the main factors that influence its clearance.

Materials and methods

This observational, single-center, cross-sectional study was performed using a cohort of monthly blood tests from the chronic hemodialysis unit. A total of 137 patients were included (47 women, 34%), with a mean age of 70 ± 17 (range: 21–96); these patients had been stable on HDF for a mean duration of 44 ± 54 months (range: 3–315). Adult patients were included in a standard schedule of 3 sessions/week of 4–5 h of duration, or in a nocturnal regimen of 8 h of duration (117 and 20, respectively). Patients with acute or unstable conditions were excluded. The underlying renal diseases included nephroangiosclerosis (16), diabetic nephropathy (17), glomerulonephritis (14), systemic disease (14), unknown (14), tubulointerstitial nephritis (12), urological nephritis (11) and renal polycystosis (3). All patients signed an informed consent form. The study was approved by the local Ethics Committee and was conducted in accordance with the Declaration of Helsinki.

All patients participated in a regular dialysis session in postdilution HDF mode with the following parameters: bicarbonate-buffered dialysate, blood flow (Qb) of 421 \pm 30 mL/min, dialysate flow (Qd) of 400 mL/min and a dialysis time (Td) of 304 \pm 56 min (range: 240–

480). The net ultrafiltration was individually configured according to the clinical needs of the patient. Vascular access was achieved using an arteriovenous fistula in 74 patients, a prosthetic arteriovenous fistula in 7 patients, and a tunneled catheter in 56 patients. Low-molecular-weight heparin (tinzaparin) was used for anticoagulation in 57.6% of the patients, sodium heparin was used in 30.7%, and the remaining 11.7% were dialyzed without anticoagulation. Fresenius 5008 CorDiax or 6008 CARE system dialysis monitors (Fresenius, Bad Homburg, Germany) were used. The following dialyzers were used: 72 FX60 (Fresenius, Bad Homburg, Germany), 39 Clearum (Bellco, Mirandola, Italy), 12 CorAL60 (Fresenius, Bad Homburg, Germany), 12 Solacea (Nipro, Osaka, Japan) and 2 FX50 (Fresenius, Bad Homburg, Germany).

The following dialysis parameters were collected: actual duration, dialyzer, Qb, recirculation index measured by the temperature module, initial and final hematocrit measured automatically using the BVM biosensor, initial and final body weights, total volume of blood processed, and replacement volume.

Blood samples were obtained from each patient for analysis. Laboratory measurements included concentrations of urea (molecular weight [MW] 60) and creatinine (MW 113) in serum at the beginning and at the end of each session to calculate the reduction rate (RR) of these solutes. Uremic toxins bound to proteins, namely, pCS (MW 108) and IS (MW 213), were also evaluated. The final IS and pCS concentration were corrected for the degree of hemoconcentration and the volume of distribution (approximate extracellular volume) according to Bergström and Wehle. ¹⁶

Urea and creatinine were measured using molecular absorption spectrometry in an Atellica Solution analyzer (Siemens Healthineers, Tarrytown, NY, USA). IS and pCS levels in serum were measured using liquid chromatography—mass spectrometry (LC–MS) according to the methodology specified in a previous study. ¹⁷

The results are expressed as the arithmetic mean \pm standard deviation. To analyze the statistical significance of the quantitative parameters, we performed a Student's t-test for independent data. To identify the factors that predict greater or lesser clearance of PBUT, univariate and multivariate logistic regression models were developed. The Pearson correlation coefficient was calculated to evaluate the linear relationship between IS RR and pCS RR and the different variables that were significant. P values < .05 were considered to indicate statistical significance. The analyses were performed using SPSS version 23 (SPSS, Chicago, IL, USA).

Results

All the sessions were performed without noteworthy incidents. The variables related to the dialysis technique are shown in Table 1. Table 1: Qb, total volume of blood processed, actual duration, vascular access recirculation index, initial weight, final weight, weight gain, initial and final hematocrit, replacement volume in HDF, total convective volume (replacement volume plus ultrafiltered volume) and the calculation of the filtration fraction (FF).

Table 1Variables related to the dialysis session.

Variable	Mean ± SD	Range		
Blood flow (mL/min)	420 ± 29.75	350-450		
Total purified blood (L)	122.29 ± 21.76	80.4-191		
Dialysis flow (mL/min)	372.26 ± 68.39	200-400		
Real time (min)	303.83 ± 55.57	230-471		
Initial weight (kg)	70.58 ± 16.19	38.1-116.6		
Final weight (kg)	68.67 ± 15.91	37.4-114.5		
Weight gain (kg)	1.91 ± 0.96	0-5		
Initial hematocrit (%)	30.42 ± 4.98	20-41.4		
Final hematocrit (%)	35.03 ± 6.15	20.8-48.7		
Recirculation (%)	13.86 ± 5.52	3-31		
Substituttion volume (L)	32.47 ± 8.76	11.4-68.6		
Total convective liters (L)	34.60 ± 8.30	13.99-61.46		
Filtration fraction (%)	28.22 ± 4.42	14.3-37.3		

^{%,} percentage; SD, standard deviation, kg, kilograms; L, liters; min, minutes; mL, milliliters.

Reduction ratios in uremic toxins

The serum concentrations of uremic toxins (BUN, creatinine, IS and pCS) and the reduction ratios are shown in Table 2. The IS RR was $53.4\pm9.3\%$ (range: 24–76), and that for pCS was $48.2\pm11.3\%$ (range: 17–76).

Association study

The greatest association was found between the RR of both PBUTs analyzed (r=0.606; P<.01). This finding shows that both toxins are eliminated by similar mechanisms.

Univariate analysis revealed that for IS, the greatest correlation occurred with the urea and creatinine RR (Fig. 1), followed by sex (Fig. 2), weight, hematocrit (Supplementary Fig. 1 in Appendix B) and the convective volume expressed as FF (Supplementary Fig. 2 in Appendix B); however, the correlation did not reach statistical

Table 2
Serum concentrations of uremic toxins.

Variable	Mean ± SD	Range
BUN pre (mg/dL)	49.47 ± 14.65	21-103
BUN post (mg/dL)	8.00 ± 3.73	4-21
RRU (%)	83.9 ± 4.9	70.7-91.8
Creatinine pre (mg/dL)	6.17 ± 1.83	2.61 - 11.78
Creatinine post (mg/dL)	1.43 ± 0.56	0.44 - 3.31
Creatinine RR (%)	76.9 ± 6.1	57.2-90.7
IS pre (ng/mL)	24,299.48 ± 13,605.96	2,989-74,521
IS post (ng/mL)	$11,158.25 \pm 6,258.66$	613-30,873
IS RR (%)	53.4 ± 9.3	24.6-75.5
pCS pre (ng/mL)	$31,936.99 \pm 19,413.39$	179-98,131
pCS post (ng/mL)	$16,469.58 \pm 10,093.52$	85-46,654
pCS RR (%)	48.2 ± 11.3	17.1-76.4

SD, standard deviation; BUN, blood urea nitrogen; pre, predialysis; post, postdialysis; mg, milligrams; dL, deciliters; ng, nanograms; mL, milliliters; RR, reduction ratio; U, urea; IS, indoxyl sulfate; pCS, p-cresyl sulfate. The final pCS and IS concentrations were corrected for the degree of hemoconcentration and the volume of distribution (approximate extracellular volume) according to Bergström and Wehle (16).

significance with albumin, dialysis time, replacement volume or total convective volume (Table 3).

Regarding pCS, the highest correlation was also observed with the urea and creatinine RR (Fig. 1), followed by sex (Fig. 2), weight, hematocrit (Supplementary Fig. 1 in Appendix B), substitution volume, total convective volume and FF (Supplementary Fig. 2 in Appendix B); the correlation with albumin or dialysis time did not reach statistical significance (Table 3). Unlike the IS RR, pCS RR shows a statistically significant correlation with the replacement volume and with the total convective volume. It also had a positive correlation with the final weight but not with the final hematocrit.

In the multivariate linear regression analysis, only 3 variables remained as independent predictors of the IS RR: sex, Td and URR




Fig. 1. Scatter plot of the percentage reduction in indoxyl sulfate with respect to the percentage reduction in urea (A), creatinine (C) and p-cresyl sulfate (B and D, respectively). The Pearson coefficients and the corresponding *P* values are shown.

V.J. Escudero-Saiz, E. Cuadrado-Payán, M. Rodríguez-García et al.

Nefrologia xx (2025) 501391




Fig. 2. Box plot of the IS RR (left) and pCS RR (right) by sex. The P values obtained using the Student's t-test are shown for each variable.

$$\begin{aligned} \text{ISRR} \, = \, \text{URR} \, + \, [(0.194 \, \times \, \text{sex}) \, + \, (-0.155 \, \times \, \text{Td}) \\ \\ + \, (0.451 \, \times \, \text{URR}) - 16.5] \end{aligned}$$

where, sex was coded as 0 for men and 1 for women, the Td is expressed in minutes, and the URR is expressed as a percentage. This model presents a multiple correlation coefficient (R) of 0.531, with a coefficient of determination (R^2) of 0.282. This indicates that 28.2% of the variability in the IS RR is explained by these 3 variables.

On the other hand, with respect to the pCS RR in the multivariate analysis, only its correlation with the creatinine RR remained statistically significant. In this case, unlike the IS RR, the creatinine RR had more statistical value than the URR did.

Discussion

The clearance of uremic toxins bound to proteins by HDF with high convective volumes in our unit is approximately 53% and 48% for IS and pCS, respectively; these results are similar to those published in the literature (Table 4). The association analysis revealed a clearance

directly proportional to the clearance of small molecules, indicating the fundamentally diffusive clearance of PBUT.

These toxins have a high degree of binding to albumin, with up to 95% binding for IS and pCS. ^{18,19} 3-Carboxy-4-methyl-5-propyl-2-furan-propanoic acid is the PBUT with the highest documented degree of binding (99%–100%). ¹⁹ These toxins bind to albumin at the Sudlow I and II binding sites; the Sudlow II site is predominant for IS and pCS. ¹⁸ This high degree of protein binding limits their clearance with conventional hemodialysis techniques despite being molecules with small molecular sizes. ¹⁵

Table 4 shows the results of different studies that have analyzed the clearance of PBUT using different hemodialysis techniques. Lowflux hemodialysis reaches a IS RR of 22% with Qb and Qd values of 200 mL/min and 300 mL/min, respectively. High-flux hemodialysis reaches RRs of approximately 33%–52% and 27%–47% for IS and pCS, respectively, in a conventional 4-h session 13,21-23 (Table 4). Predilution HDF achieves a IS RR of 48% and pCS RR of 41%–45%, 24,25 whereas postdilution HDF achieves a similar RR or a RR with discrete increases of 45%–55% and 38%–48% for IS and pCS, respectively. 11,13,23,26,27 As time increases (8 h), these techniques achieve RR values of 60% and 52% for IS and pCS, respectively. In addition, postdilution HDF values of 43%–55% and 37%–45% are obtained for IS and pCS, respectively, with high-flux hemodialysis 13,28

Table 3Correlation analysis.

	Indoxyl sulfate			p-Cresyl sulfate				
	Univariate		Multivariate		Univariate		Multivariate	
	Pearson Correlation	P-value	Beta coefficient	P-value	Pearson Correlation	P-value	Beta coefficient	P value
Sex	0.337	<.001	0.194	.014	0.227	.008		
Dialysis time (min)	-0.042	.629	-0.155	.047	0.066	.446		
Weight (kg)	-0.259	.002			0.235	.006		
Replacement volume (L)	0.155	.071			0.19	.027		
Total convective volume (L)	0.14	.105			0.184	.032		
Filtration fraction (%)	0.17	.048			0.178	.038		
Hematocrit (%)	-0.185	.03			-0.165	.057		
Urea RR (%)	0.466	<.001	0.451	<.001	0.328	<.001		
Creatinine RR (%)	0.442	<.001			0.383	<.001	0.383	<.001

^{%,} percentage; kg, kilograms; L, liters; Min, minutes; RR, reduction ratio.

V.J. Escudero-Saiz, E. Cuadrado-Payán, M. Rodríguez-García et al.

Nefrologia xx (2025) 501391

Table 4Percentages of reduction in uremic toxins bound to proteins reported in the main published studies.

Study	N	Technique	Qb (mL/min)	Qd (mL/min)	Time (min)	IS RR (%)	pCS RR (%)
		Post-HDF	311	619	248	44.8	40.4
Meert (2009)	14	Pre-HDF	315	535	249	48.5	41.9
		Pre-HF	311	0	251	33.8	30.6
		HD high flow (PU-)	378 ± 33	500	229 ± 22	50.4 ± 2.6	45.6 ± 2
Krieter (2010)	8	HD high flow (PU+)	378 ± 33	500	229 ± 22	52.2 ± 12.2	47.3 ± 14.8
Krieter (2010)	0	HDF (PU-)	378 ± 33	500	229 ± 22	53.3 ± 8.9	40.4 ± 25.3
		HDF (PU+)	378 ± 33	500	229 ± 22	54.6 ± 8.7	47.8 ± 10.3
		HD – BF	250	500	240	42.4	32.9
Lesafler (2010)	10	HD high flow (FX60)	250	500	240	42.6	41.4
		HD high flow (CTA)	250	500	240	41.5	32.4
Meijers (2011)	32	HD high flow (FX80)	225 ± 19	300	471 ± 16	43	37
14 (0011)	1.4	Pre-HDF	300	800	251 ± 19	48	45
Meert (2011)	14	Post HDF	300	800	252 ± 19	48	45
	_	HD high flow (high KoA)	270	800	473 ± 20	66 ± 6	59 ± 8
Sirich (2012)	9	HD high flow (low KoA)	350	300	473 ± 20	46 ± 9	41 ± 11
mt	14	OL-HDF	343	725	249	48.7	44
Eloot (2012)		Mild-dilution HDF	343	650	251	47.4	42.7
Brettschneider (2013)	5	PFD	226 ± 3	NA	300	78	71
		HD high flow	300	600	246 ± 4	35	30
		HD high flow	300	600	486 ± 2	55	45
Cornelis (2015)	13	OL-HDF	300	600	245 ± 3	45	38
		OL-HDF	300	600	487 ± 6	60	52
Abad (2016)	14	OL-HDF	426	NA	240	47.8	44.4
		HFR	>350	NA	240	48.8	50.7
Esquivias-Motta (2017)	17	HD high flow	>350	NA	240	43.3	42.2
1		HDF	>350	NA	240	45.2	39.8
Paats (2020)		Standard (HD/HDF)	323	458	240	NA	NA
	78	HD low flow	200	301	240	22	NA
		Medium HDF	306	793	240	42	NA
		High HDF	378	793	240	48	NA
Chen (2020)	37	HFR	249 ± 19	500	240	43.6	40.9

FPAD, fractionated plasma separation and adsorption technique; HD, hemodialysis; OL-HDF, online hemodiafiltration; Pre-HF, predilution hemofiltration; HFR, hemodiafiltration with endogenous reinfusion; IS, indoxyl sulfate; min, minutes; mL, milliliters; n, number; NA, not available; pCS, p-cresyl sulfate; RR, reduction ratio; pre-HDF, predilution hemodiafiltration; post-HDF, postdilutional hemodiafiltration; PU, PUREMA membrane; Qb, blood flow; Qd, dialysis flow; CTA, cellulose triacetate.

and up to 66% and 59%, respectively, with high-flux hemodialysis with high-KoA membranes¹⁴ because of the constant dissociation and the greater availability of the free fraction of the PBUT. 13 The disparity of results in the literature could be explained by the heterogeneity of variables that influence the clearance of small molecules, namely, Qb, Qd and Td, as well as by the small number of patients studied, which limits their external validity (Table 4). Expanded hemodialysis does not achieve higher RRs than previous techniques. 12 In this study, the RR values of IS and pCS are 53% and 48%, respectively, for HDF with high convective volume. The results are consistent with published works (Table 4). In our case, the nocturnal cohort did not reach a higher RR than the daytime cohort. This difference may be due to the lower Od prescribed (200 mL/min) compared with that reported by Cornelis et al.¹³ and Sirich et al.,¹ which had Qd values of 600 mL/min and 800 mL/min, respectively, as well as the difference between the two cohorts in our study (20 vs. 117). More recently, our group evaluated the possible hemoadsorption added value for the clearance of various uremic toxins using various hemodialysis modalities. Among the modalities, HDF achieved the greatest reduction in IS (55%) and pCS (51%) compared with low- or high-flux hemodialysis (a reduction that was approximately 5% lower); however, no differences in the clearance of PBUT were observed when the hemoadsorption cartridge was added in any of the 3 treatment modalities studied.²⁹

The correlation analysis revealed a relationship between the clearance of IS and pCS and the clearance of small-molecular-weight uremic toxins because diffusion is the main mechanism of their elimination^{10,13} (Fig. 1). In this sense, increasing the variables that increase the diffusive capacity of the hemodialysis technique, such as Qb, Qd, Td, or frequency of therapy, increases the clearance of PBUT in a directly proportional way. In addition, the differences in PBUT clearance between men and women (Fig. 2 and Supplementary Fig.

1 in Appendix B) are inversely proportional to the volume of distribution, since it is generally lower in females. 30 Postdilution HDF increases PBUT clearance by the discrete increase it provides in the clearance of small molecules (URR and creatinine RR) rather than by the convective effect, 31,32 as previously observed by Abad et al. 26 (Supplementary Fig. 2 in Appendix B).

Currently, different strategies are being developed to increase the clearance of PBUT. The use of competing molecules during dialysis sessions seems to be the most common in the clinical setting.³³ Specifically, the use of tryptophan, furosemide and ibuprofen has been demonstrated in several in vitro studies,34 the latter of which demonstrated the greatest competitive capacity for IS and pCS. 35 The use of salvianolic acids³⁶ and lipid emulsions (Intralipid™, Fresenius KABI SSPC, Jiangsu, China) have also been studied in murine models³⁷; however, no data from humans or on its possible safety have been reported to date.³⁸ The only competitor used clinically in patients was ibuprofen,³⁹ and clearance increased from 6 to 20.2 mL/min and from 4.4 to 14.9 for IS and pCS, respectively. Our group showed increases of 14.2% and 12.9% in the IS and pCS RR, respectively, after arterial line infusion for one hour, reaching RRs of 58.8% and 54.6%, respectively. 40 Another strategy is the development of dialysis membranes capable of adsorbing PBUT by incorporating adsorbent particles such as activated carbon, 38 zeolites 41 or metalloorganic zirconium structures. 42 In addition, the combination of both strategies with membranes containing competing molecules adhered to the luminal surface has been proposed. 43 However, no clinical data on adsorptive strategies are available to date.

Another strategy reported in the literature to reduce the PBUT concentration is dietary intervention or the use of oral adsorbents. 44 With respect to dietary modification, the MEDIKA study, a prospective crossover study conducted in patients with advanced chronic kidney disease, revealed that a very-low-protein diet (protein concentration

V.J. Escudero-Saiz, E. Cuadrado-Payán, M. Rodríguez-García et al.

Nefrologia xx (2025) 501391

of 0.3–0.5 g/kg per day, along with that of keto analogs), followed by a Mediterranean diet (protein concentration of 0.7–0.8 g/kg per day with a predominance of vegetable origin), resulted in reduced IS and pCS concentrations compared with a standard diet (1 g/kg per day with a predominance of animal origin).⁴⁵ The authors explain this effect using a double mechanism. The first mechanism involves a reduction in protein intake, which is the initial substrate of proteinbound uremic toxins, and the second mechanism involves a modulatory effect on the intestinal microbiota, with a reduction in proteobacteria and an increase in saccharolytic and butyrate-forming species. 43 With respect to the use of oral adsorbents as chelators of the products of the metabolism of the intestinal microbiota, there are only data on the use of Kremezin® (spherical activated carbon, AST-120; Kureha Chemical Industry Co Ltd, Tokyo, Japan) in patients with advanced chronic kidney disease, 46 in whom a dose-dependent reduction in IS was observed. In addition, in hemodialysis patients, ⁴⁷ the total IS and total pCS levels were reduced by 45% and 31%, respectively.

This study has several notable limitations. First, the absence of a control group with another modality of hemodialysis prevents a direct comparison of the effect of postdilution HDF. Second, despite the influence of the duration of dialysis on the diffusion capacity of these molecules, the large difference between the 2 cohorts presented (117 vs. 20 patients) could explain the lack of statistically significant differences, as well as the discrepancies with the results of other studies. Third, only the clearance of 2 PBUT (IS and pCS) was measured, which makes it difficult to generalize to other uremic toxins bound to proteins. Finally, although the multivariate model explains part of the variability in IS clearance, other factors not considered could potentially influence the results. However, the main strength of this work is that it represents the largest cohort studied on the clearance of PBUT. In addition, our results reinforce the idea of the use of high blood and dialysate flows, along with an increase in therapy duration within each unit, to improve the diffusive capacity of these uremic toxins.

In summary, postdilution HDF with high convective volumes achieves clearance rates of protein-bound uremic toxins similar to those reported by other groups but higher than those obtained with high-flux hemodialysis or expanded hemodialysis. However, these results are insufficient given the demonstrated toxicity of these toxins, which cause increased cardiovascular mortality in hemodialysis patients. To optimize clearance, future studies should evaluate combined strategies, such as the integration of adsorptive membranes with hemodialysis techniques or the use of competing molecules.

Financing

This research has not received specific support from public sector agencies, the commercial sector or nonprofit entities.

Declaration of competing interest

The authors declare that they have not received financial support for this project. FM has received fees from Amgen, Baxter, Fresenius Medical Care, Medtronic, Nipro, Palex and Vifor. The remaining authors declare that they have no conflicts of interest.

Acknowledgments

We would like to express our gratitude to all the patients who participated and to all the staff of the Dialysis Section of the Hospital Clínic de Barcelona for their collaboration in this study.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:https://doi.org/10.1016/j.nefroe.2025. 501391

References

- Rosner MH, Reis T, Husain-Syed F, Vanholder R, Hutchison C, Stenvinkel P, et al. Classification of uremic toxins and their role in kidney failure. Clin J Am Soc Nephrol. 2021;16:1918–28.
- European Uremic Toxins Work Group. List of uremic solutes—uremic solutes database. [Internet] [accessed 28 Oct 2024]. Available from: https://database. uremic-toxins.org/soluteList.php.
- Vanholder R, Pletinck A, Schepers E, Glorieux G. Biochemical and clinical impact of organic uremic retention solutes: a comprehensive update. Toxins (Basel). 2018;10:1–57.
- Dehghan Niestanak V, Unsworth LD. Detailing protein-bound uremic toxin interaction mechanisms with human serum albumin in the pursuit of designing competitive binders. Int J Mol Sci. 2023;24.
- Daneshamouz S, Eduok U, Abdelrasoul A, Shoker A. Protein-bound uremic toxins (PBUTs) in chronic kidney disease (CKD) patients: production pathway, challenges and recent advances in renal PBUTs clearance. NanoImpact [Internet]. 202121100299, http://dx.doi.org/10.1016/j.impact.2021.100299 (January).
- Wu IW, Hsu KH, Hsu HJ, Lee CC, Sun CY, Tsai CJ, et al. Serum free p-cresyl sulfate levels predict cardiovascular and all-cause mortality in elderly hemodialysis patients—a prospective cohort study. Nephrol Dial Transplant. 2012;27:1169–75.
- Bammens B, Evenepoel P, Keuleers H, Verbeke K, Vanrenterghem Y. Free serum concentrations of the protein-bound retention solute p-cresol predict mortality in hemodialysis patients. Kidney Int [Internet]. 2006;69:1081–7, http://dx.doi.org/ 10.1038/si.ki.5000115
- Meijers BKI, Bammens B, De Moor B, Verbeke K, Vanrenterghem Y, Evenepoel P. Free p-cresol is associated with cardiovascular disease in hemodialysis patients. Kidney Int [Internet]. 2008;73:1174–80, http://dx.doi.org/10.1038/ki.2008.31
- Wanner C, Vanholder R, Ortiz A, Davenport A, Canaud B, Blankestijn PJ, et al. Proceedings of a membrane update symposium: advancements, scientific insights, and future trends for dialysis membranes for enhanced clinical outcomes in end stage kidney disease patients. Front Nephrol. 2024;1–12.
- Saar-Kovrov V, Zidek W, Orth-Alampour S, Fliser D, Jankowski V, Biessen EAL, et al. Reduction of protein-bound uraemic toxins in plasma of chronic renal failure patients: a systematic review. J Intern Med. 2021;290:499–526.
- Krieter DH, Hackl A, Rodriguez A, Chenine L, Moragues HL, Lemke HD, et al. Protein-bound uraemic toxin removal in haemodialysis and post-dilution haemodiafiltration. Nephrol Dial Transplant. 2010;25:212–8.
- Kim YG, Lee SH, Jung SW, Jung GT, Lim HJ, Kim KP, et al. The medium cut-off membrane does not lower protein-bound uremic toxins. Toxins (Basel). 2022;14 (779).
- 13. Cornelis T, Eloot S, Vanholder R, Glorieux G, Van Der Sande FM, Scheijen JL, et al. Protein-bound uraemic toxins, dicarbonyl stress and advanced glycation end products in conventional and extended haemodialysis and haemodiafiltration. Nephrol Dial Transplant. 2015;30:1395–402.
- Sirich TL, Luo FJG, Plummer NS, Hostetter TH, Meyer TW. Selectively increasing the clearance of protein-bound uremic solutes. Nephrol Dial Transplant. 2012;27:1574–9.
- Sánchez-Ospina D, Mas-Fontao S, Gracia-Iguacel C, Avello A, González de Rivera M, Mujika-Marticorena M, et al. Displacing the burden: a review of protein-bound uremic toxin clearance strategies in chronic kidney disease. J Clin Med. 2024;13(5).
- Bergström J, Wehle B. No change in corrected β2-microglobulin concentration after cuprophane haemodialysis. Available from: Lancet [Internet]. 1987;329:628–9, https://www.sciencedirect.com/science/article/pii/S0140673687902662.
- Rodríguez-García M, Martínez I, Aliart I, Sainz de Medrano JI, Rico N, Escudero-Saiz VJ, et al. Validation of an LC-HRMS method for quantifying indoxyl sulfate and p-cresyl sulfate in human serum. Available from: Molecules [Internet]. 2025;30:782. https://www.mdni.com/1420-3049/30/4/782.
- Watanabe H, Noguchi T, Miyamoto Y, Kadowaki D, Kotani S, Nakajima M, et al. Interaction between two sulfate-conjugated uremic toxins, p-cresyl sulfate and indoxyl sulfate, during binding with human serum albumin. Available from: Drug Metabol Dispos [Internet]. 2012;40:1423–8, http://dmd.aspetjournals.org/ content/40/7/1423. Abstract.
- Shi Y, Tian H, Wang Y, Shen Y, Zhu Q, Ding F. Effect of ionic strength, pH and chemical displacers on the percentage protein binding of protein-bound uremic toxins. Blood Purif. 2019;47:351–60.
- 20. Paats J, Adoberg A, Arund J, Dhondt A, Fernström A, Fridolin I, et al. Serum levels and removal by haemodialysis and haemodiafiltration of tryptophan-derived uremic toxins in ESKD patients. Int J Mol Sci. 2020;21(4).
- Lesaffer G, De Smet R, Lameire N, Dhondt A, Duym P, Vanholder R. Intradialytic removal of protein-bound uraemic toxins: role of solute characteristics and of dialyser membrane. Nephrol Dial Transplant. 2000;15:50–7.
- Luo FJG, Patel KP, Marquez IO, Plummer NS, Hostetter TH, Meyer TW. Effect of increasing dialyzer mass transfer area coefficient and dialysate flow on clearance of protein-bound solutes: a pilot crossover trial. Am J Kidney Dis. 2009;53:1042–9.
- 23. Esquivias-Motta E, Martín-Malo A, Buendia P, Álvarez-Lara MA, Soriano S, Crespo R, et al. Hemodiafiltration with endogenous reinfusion improved

V.J. Escudero-Saiz, E. Cuadrado-Payán, M. Rodríguez-García et al.

microinflammation and endothelial damage compared with online-

- hemodiafiltration: a hypothesis generating study. Artif Organs. 2017;41:88–98.
 Meert N, Eloot S, Waterloos MA, Van Landschoot M, Dhondt A, Glorieux G, et al. Effective removal of protein-bound uraemic solutes by different convective strategies: a prospective trial. Nephrol Dial Transplant. 2009;24:562–70.
- 25. Meert N, Eloot S, Schepers E, Lemke HD, Dhondt A, Glorieux G, et al. Comparison of removal capacity of two consecutive generations of high-flux dialysers during different treatment modalities. Nephrol Dial Transplant. 2011;26:2624–30.
- Abad S, Vega A, Quiroga B, Arroyo D, Panizo N, Reque JE, et al. Toxinas unidas a proteínas: valor añadido en su eliminación con altos volúmenes convectivos. Nefrología. 2016;36:637–42.
- Eloot S, Dhondt A, Van Landschoot M, Waterloos MA, Vanholder R. Removal of water-soluble and protein-bound solutes with reversed mid-dilution versus postdilution haemodiafiltration. Nephrol Dial Transplant. 2012;27:3278–83.
- Meijers B, Toussaint ND, Meyer T, Bammens B, Verbeke K, Vanrenterghem Y, et al. Reduction in protein-bound solutes unacceptable as marker of dialysis efficacy during alternate-night nocturnal hemodialysis. Am J Nephrol. 2011;34:226–32.
- Maduell F, Escudero-Saiz VJ, Cuadrado-Payán E, Rodriguez-Garcia M, Rodas LM, Fontseré N, et al. Comparing hemodialysis and hemodiafiltration performance with and without hemoadsorption. Clin Kidney J. 2025;18(5).
- Maduell F, Miralles F, Caridad A, Singüenza F, Serrato F, Ocho E. Análisis del volumen de distribución de la urea en hemodiálisis. Available from: Nefrologia [Internet]. 1992;12:411–5, http://www.elsevier.es,day31/01/.
- Maduell F, Del Pozo C, Garcia H, Sanchez L, Hdez-Jaras J, Albero MD, et al. Change from conventional haemodiafiltration to on-line haemodiafiltration. Nephrol Dial Transplant. 1999;14.
- Lin CL, Huang CC, Yu CC, Wu CH, Chang CT, Hsu HH, et al. Improved iron utilization and reduced erythropoietin resistance by on-line hemodiafiltration. Blood Purif. 2002:20.
- Maheshwari V, Tao X, Thijssen S, Kotanko P. Removal of protein-bound uremic toxins using binding competitors in hemodialysis: a narrative review. Toxins (Rasel) 2021:13
- 34. Tao X, Thijssen S, Kotanko P, Ho CH, Henrie M, Stroup E, et al. Improved dialytic removal of protein-bound uraemic toxins with use of albumin binding competitors: an in vitro human whole blood study. Sci Rep. 2016;6:2–10.
- Maheshwari V, Thijssen S, Tao X, Fuertinger DH, Kappel F, Kotanko P. In silico comparison of protein-bound uremic toxin removal by hemodialysis,

hemodiafiltration, membrane adsorption, and binding competition. Sci Rep [Internet]. 2019;9:1–13, http://dx.doi.org/10.1038/s41598-018-37195-1

Nefrologia xx (2025) 501391

- Li J, Wang Y, Xu X, Cao W, Shen Z, Wang N, et al. Improved dialysis removal of protein-bound uremic toxins by salvianolic acids. Phytomedicine. 2019;57:166–73.
- Shi Y, Zhang Y, Tian H, Wang Y, Shen Y, Zhu Q, et al. Improved dialytic removal of protein-bound uremic toxins by intravenous lipid emulsion in chronic kidney disease rats. Nephrol Dial Transplant. 2019;34:1842–52.
- **38.** Rodrigues FSC, Faria M. Adsorption- and displacement-based approaches for the removal of protein-bound uremic toxins. Toxins (Basel). 2023;15(110).
- Madero M, Cano KB, Campos I, Tao X, Maheshwari V, Brown J, et al. Removal of protein-bound uremic toxins during hemodialysis using a binding competitor. Clin J Am Soc Nephrol. 2019:14:394–402.
- 40. Escudero-Saiz VJ, Cuadrado-Payán E, Rodriguez-Garcia M, Casals G, Rodas LM, Fontseré N, et al. The choice of anti-inflammatory influences the elimination of protein-bound uremic toxins. Toxins. 2024;16(Dec (12)).
- Lu L, Yeow JTW. An adsorption study of indoxyl sulfate by zeolites and polyethersulfone–zeolite composite membranes. Mater Des. 2017;120:328–35.
- Zeng S, Hou Y, Zhou Y, Zhou X, Ye S, Wang M, et al. Adsorptive removal of uremic toxins using Zr-based MOFs for potential hemodialysis membranes. J Mater Sci. 2022;57:2909–23.
- Rodrigues FSC, Brilhante D, Macêdo A, Pires RF, Faria M. Ibuprofen-immobilized thin films: a novel approach to improve the clearance of protein-bound uremic toxins. ACS Appl Mater Interfaces. 2024;16:6589–604.
- 44. Vanholder R, Snauwaert E, Verbeke F. Future of uremic toxin management;2024;1–
- 45. Di Iorio BR, Rocchetti MT, De Angelis M, Cosola C, Marzocco S, Di Micco L, et al. Nutritional therapy modulates intestinal microbiota and reduces serum levels of total and free indoxyl sulfate and p-cresyl sulfate in chronic kidney disease (Medika study). J Clin Med. 2019;8(9).
- Schulman G, Agarwal R, Acharya M, Berl T, Blumenthal S, Kopyt N. A multicenter, randomized, double-blind, placebo-controlled, dose-ranging study of AST-120 (Kremezin) in patients with moderate to severe CKD. Am J Kidney Dis. 2006:47:565–77
- Yamamoto S, Kazama JJ, Omori K, Matsuo K, Takahashi Y, Kawamura K, et al. Continuous reduction of protein-bound uraemic toxins with improved oxidative stress by using the oral charcoal adsorbent AST-120 in haemodialysis patients. Sci Rep. 2015;5