



Revista de la Sociedad Española de Nefrología

journal homepage: www.revistanefrologia.com



Original article

Q1 Analysis of the association between Healthy Eating Index (HEI)-2020 and diabetic kidney disease in type 2 diabetes patients: Based on the NHANES database

Q2 Chunping Dong, Hui Li, Jing Li, Yuan Qiao, Shan Gao*

Department of Endocrinology, Shaanxi Provincial People's Hospital, Xi'an City 710068, Shaanxi Province, China

ARTICLE INFO

Keywords:

Type 2 diabetes
Diabetic kidney disease
HEI-2020
National Health and Nutrition Examination Survey

ABSTRACT

Background: A rise in the Healthy Eating Index (HEI) 2020 is associated with a reduced risk of chronic kidney disease. However, it remains unclear whether HEI-2020 exhibits a similar positive effect in diabetic patients. **Objectives:** This work aimed to investigate the relationship between HEI-2020 and the prevalence of diabetic kidney disease (DKD) in type 2 diabetes (T2D) patients.

Methods: A cross-sectional analysis was based on the National Health and Nutrition Survey database from 2007 to 2018. A weighted logistic regression model, restricted cubic splines, and subgroup analysis were undertaken to assess the relationship between HEI-2020 and the prevalence of DKD in T2D patients. A weighted logistic regression model and a weighted quantile sum (WQS) regression model were employed. The independent impact of HEI-2020-related components on the prevalence of DKD among T2D patients was analyzed. In addition, sensitivity analysis was conducted.

Results: A total of 4255 participants with T2D were enrolled, with a DKD prevalence rate of 34.7%. The logistics regression analysis demonstrated a significant inverse relation between the HEI-2020 score and the prevalence of DKD in fully adjusted models (OR = 0.982, 95% CI: 0.973–0.992, $P < 0.001$). No nonlinear relationship was detected (P -non-linear = 0.970). No subgroup with significant interaction was identified. The WQS regression indicated that in fully adjusted models, the WQS index for HEI-2020 was significantly negatively correlated with the prevalence of DKD among T2D patients (OR = 0.765, 95% CI: 0.621–0.943, $P = 0.012$), with dairy products, total protein foods, and total vegetables as the most influential components. In addition, HEI-2020 was significantly negatively correlated with DKD in previously diagnosed T2D patients (OR = 0.979, 95% CI: 0.967–0.990, $P < 0.001$), while no significant association was observed in newly diagnosed patients. The sensitivity analysis yielded similar results, indicating that our results were robust. **Conclusion:** This research revealed a relationship between an increase in the HEI-2020 score and a lower prevalence of DKD among T2D patients, emphasizing the importance of the intake of dairy products, total protein foods, and total vegetables. This finding provides new scientific grounds for dietary management in T2D patients.

Introduction

Q3 A total of 589 million patients aged 20–79 are diagnosed with diabetes in the world. Type 2 diabetes (T2D) is the most prevalent form of diabetes globally, accounting for 96.0% of all diabetes cases, and is associated with numerous complications that significantly impact patient quality of life and life expectancy.^{1,2} Diabetic kidney disease (DKD), a key microvascular complication of T2D, affects roughly 50% of T2D patients who eventually progress to this condition.³ In 2021, the global number of DKD cases related to

T2D will reach 107 million, with 477,000 deaths. The age-standardized incidence rate is expected to continue to rise in the future,⁴ placing an immense burden on global public health. Moreover, DKD is associated with excess mortality in diabetes patients, with the crude all-cause mortality rate in the DKD cohort being nearly twice that of the control group (simple T2D patients) (50.3 vs 28.4 per 1000 person-years).⁵ DKD remains a primary cause of end-stage renal disease, triggering arterial diseases and cardiovascular complications that complicate treatment for patients.⁶ Consequently, identifying means to curb the progression of DKD is crucial for reducing the mortality risk among T2D patients.

Nutrition interventions are key to diabetes management, with a balanced diet contributing to improved glucose metabolism and

* Corresponding author.

E-mail address: gaoss4762367@163.com (S. Gao).

<https://doi.org/10.1016/j.nefro.2025.501452>

Received 14 July 2025; Accepted 11 November 2025

Available online xxx

0211-6995/© 2025 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

reduced DKD risk in patients with diabetes.⁷ The United States Dietary Guidelines recommend assessing the quality of American diets using the Healthy Eating Index (HEI), which serves to identify healthy dietary patterns for various populations. The HEI is updated in tandem with the Dietary Guidelines, with the most recent version being HEI-2020.⁸ It has been widely employed in research on chronic diseases, with the highest quintile array of HEI scores demonstrating significantly lower risk for chronic diseases (including diabetes) compared to the lowest quintile array.⁹ A systematic review has also shown that better adherence to HEI can lower risks for diabetes and cardiovascular diseases (CVD).¹⁰ It is noteworthy that higher HEI adherence can lower the risk of CVD in diabetes patients by 50%.¹¹ Given this, we speculate that enhancing HEI adherence is likely to also lower the risk of DKD in patients with diabetes. Moreover, there is evidence that an increase in HEI-2020 scores is linked with a reduced risk of chronic kidney disease (CKD),¹² although the active effect on diabetic patients has not yet been adequately validated.

Thus, this work investigates the relationship between HEI-2020 scores and the prevalence of DKD in T2D patients enrolled in the National Health and Nutrition Examination Survey (NHANES) database, aiming to shed valuable insights into the prevention and dietary guidance for T2D patients with DKD.

Methods

Study population

The population data in this project were available through the NHANES database (<http://www.cdc.gov/nchs/nhanes.htm>), a nationally representative health survey conducted biennially with approval from the National Center for Health Statistics of the US Centers for Disease Control and Prevention. The database is used to evaluate the health and nutrition status of US adults and children.

An analysis of data spanning six cycles from 2007 to 2018 ($n = 59,842$) was conducted. Participants with diabetes were identified by the following criteria: (1) fasting blood glucose concentration ≥ 126 mg/dL, or (2) glycated hemoglobin A1c (HbA1c) level $\geq 6.5\%$, or (3) 2-h blood glucose ≥ 200 mg/dL during the 75 g oral glucose tolerance test (OGTT), or (4) self-reported diabetes or insulin or oral hypoglycemic drug use.¹³ Participants without diabetes ($n = 48,903$), missing and useless DKD data ($n = 2077$), and other missing or unavailable covariate data ($n = 4607$) were excluded. Subsequently, we screened and excluded the possible type 1 diabetes patients according to the criteria of diagnosis age < 30 years and insulin treatment. In this study, no patients ($n = 0$) met criteria.¹⁴ Finally, a total of 4255 T2D participants were included (Fig. 1). Furthermore, according to the diagnosis time of T2D, the study population was divided into a previous diagnosis group (self-reported diabetes or using hypoglycemic drugs) and a new diagnosis group (no previous diagnosis or medication history, but fasting blood glucose concentration ≥ 126 mg/dL or HbA1c $\geq 6.5\%$ or OGTT 2-h blood glucose ≥ 200 mg/dL).

The measurement of HEI-2020

Food intake data were collected via 24-h dietary recall, including participants with two complete 24-h dietary recalls that reliably met the minimum standards. HEI-2020 is the latest version used to assess dietary quality, with its components identical to those of HEI-2015.⁸ These components are divided into two forms: nine adequate components (total fruits, total vegetables, whole fruits, vegetables and legumes, dairy products, total protein foods, whole grains, seafood and plant proteins, and fatty acids), and four moderate components (refined grains, added sugars, sodium, and saturated fats). The higher intake of adequate components indicated the higher scores. On the other hand, moderate components were the opposite,




Fig. 1. Diagram of inclusion and exclusion.

with higher intake resulting in lower scores.¹⁵ The highest scores for total fruits, whole fruits, total vegetables, vegetables and legumes, total protein foods, and seafood and plant-based proteins are 5. The maximum score for the remaining seven components is 10. The HEI-2020 score ranges from 0 to 100, with higher scores reflecting higher dietary quality.

Definition of diseases

The kidney diseases were diagnosed based on the Kidney Disease: Improving Global Outcome (KDIGO). DKD was defined as diabetes combined with proteinuria (urinary albumin/creatinine ratio ≥ 30 mg/g) and/or decreased estimated glomerular filtration rate (eGFR) (eGFR < 60 mL/min/1.73 m²).¹⁶ The calculation of eGFR was based on the CKD Epidemiology Collaboration equation for standardized creatinine¹⁷ (eGFR = $141 \times \min(\text{Scr}/\kappa, 1)^\alpha \times \max(\text{Scr}/\kappa, 1)^{-1.209} \times 0.993^{\text{Age}} \times 1.018$ [if female] $\times 1.159$ [if black]; Scr: creatinine concentration; κ values: assigned according to gender, 0.9 for males and 0.7 for females; α values: determined by gender, -0.411 for males and -0.329 for females).

Covariates

The covariates included demographic characteristics, lifestyle factors, medical history, and laboratory indicators (Table S1).

Statistical analysis

All statistical analyses were performed in R (version 4.4.1). Baseline characteristics were summarized using the “tableone” package, which facilitated automated descriptive tables (sample size and percentage (n (%)) represented categorical variables; mean and standard deviation (mean (SD)) represented continuous variables). We employed the “survey” package to set up a weighted logistics regression model to elucidate the association between HEI-2020 scores and the prevalence rate of DKD in T2D patients, with trend analysis conducted. This study used the “rms” (regression modeling strategies) package to construct the restricted cubic spline and set nodes at the 5th, 50th, and 95th percentiles of the HEI-2020 score to evaluate and explore the nonlinear relationship between the HEI-2020 score and the prevalence of DKD in patients with T2D. Subgroup analyses were undertaken to examine these associations in different subpopulations.

Furthermore, a weighted logistic regression model was employed to determine the relation between 13 independent components of HEI-2020 and the prevalence of DKD in T2D patients. A weighted quantile sum (WQS) regression model was built to evaluate the contributions of 13 independent components related to HEI-2020. In the WQS model, all component effects were constrained in the same direction, with their total weights summing up to 1.¹⁸ We estimated the individual weights for each component using the quintile arrays ($q = 5$) in HEI-2021. To estimate the parameters, 60% of the data were randomly classified into a training dataset, and the remaining 40% was assigned to a validation dataset. Furthermore, considering the impact of the duration of T2D on DKD, we explored the relationship between HEI-2020 and DKD in the previously diagnosed group and the newly diagnosed group, respectively. Since some drugs, including angiotensin converting enzyme inhibitors (ACEIs), angiotensin II receptor antagonists (ARBs), sodium glucose cotransporter 2 (SGLT2) inhibitors, glucagon-like peptide-1 (GLP-1) receptor agonists, and statins, have been shown to be beneficial for renal outcomes,¹⁹ we adjusted for these factors before conducting further sensitivity analysis.

$P < 0.05$ indicated statistical significance.

Results

Baseline characteristics

A total of 4255 participants with T2D were analyzed, with 1600 diagnosed with DKD, representing a prevalence of 34.7%. The baseline characteristics of the study sample included an average age of 59.22 ± 13.37 years across the entire population, with approximately equal proportions of males and females (Table 1). Compared with patients without DKD, those with DKD exhibited lower HEI-2020 scores (51.13 ± 11.32 vs. 52.33 ± 11.72 , $P = 0.036$). Moreover, DKD patients were older, had a history of smoking, hypertension, intense physical activity (PA), a larger waist circumference, higher levels of blood urea nitrogen and uric acid, without drinking habits, and with lower high-density lipoprotein levels (all P -values < 0.05).

The relationship between the HEI-2020 score and DKD prevalence in T2D patients

To probe into the relationship between the HEI-2020 score and DKD prevalence in T2D patients, a weighted logistic regression model was constructed. Results from the fully adjusted model revealed a significant negative association between HEI-2020 scores and the prevalence of DKD in T2D patients (OR = 0.982, 95% CI: 0.973–0.992, $P < 0.001$) (Table 2). In both adjusted models, with each increase in the tertile range, the risk of DKD among T2D patients exhibited a trend towards decreased risk (P for trend < 0.001). Additionally, in Model 2, the prevalence of DKD among T2D patients in the third tertile array was significantly lower than that in the first tertile array (OR = 0.666, 95% CI: 0.518–0.857, 0.002) (Table 2).

Restricted cubic spline revealed a significant overall trend between HEI-2020 scores and the prevalence of DKD in T2D patients (P -overall < 0.001), but no nonlinear relationship was detected (P -nonlinear = 0.970) (Fig. 2).

Subgroup analysis

The subgroup analysis revealed that HEI-2020 scores were negatively correlated with the prevalence of DKD in patients with T2D for those with body mass index (BMI) between 25 and 30 kg/m² (OR = 0.977, 95% CI: 0.960–0.994, $P = 0.008$) or ≥ 30 kg/m² (OR = 0.986, 95% CI: 0.973–0.998, $P = 0.025$), those without smoking habit (OR = 0.977, 95% CI: 0.963–0.991, $P = 0.001$), those previously smoked (OR = 0.981, 95% CI: 0.963–0.999, $P = 0.034$), individuals with hypertension (OR = 0.982, 95% CI: 0.971–0.992, $P = 0.001$), and those with intense physical activity (OR = 0.977, 95% CI: 0.962–0.992, $P = 0.003$). Furthermore, no significant interactions were detected among the subgroups (P for interaction > 0.05) (Fig. 3).

The relationship between the HEI-2020 components and DKD prevalence in T2D patients

A weighted logistic regression model was constructed to examine the independent effects of 13 dietary components on the HEI-2020. OR values were all less than 1 for all components, with a significant negative correlation detected between DKD prevalence among T2D and components such as total vegetable intake (OR = 0.916, 95% CI: 0.841–0.999, $P = 0.043$), dairy (OR = 0.953, 95% CI: 0.911–0.997, $P = 0.032$), total protein foods (OR = 0.884, 95% CI: 0.796–0.981, $P = 0.018$), seafood and plant proteins (OR = 0.902, 95% CI: 0.839–0.970, $P = 0.005$), added sugars (OR = 0.937, 95% CI: 0.888–0.989, $P = 0.017$) (Table 3).

The WQS regression model was further utilized to assess the impact of HEI-2020-related diet component intakes on the prevalence of DKD

Table 1
Characteristics of NHANES type 2 diabetes participants between 2007 and 2018.

Characters	Total	Non-DKD	DKD	P-value
Overall	4255	2655 (65.3)	1600 (34.7)	
Age	59.22 (13.37)	56.72 (12.85)	63.92 (13.07)	<0.001
Race				0.253
Mexican American	745 (10.0)	485 (10.2)	260 (9.7)	
Other Hispanic	470 (6.0)	314 (6.7)	156 (4.7)	
Non-Hispanic White	1614 (62.4)	941 (61.7)	673 (63.7)	
Non-Hispanic Black	1054 (13.9)	666 (13.6)	388 (14.3)	
Other race	372 (7.8)	249 (7.8)	123 (7.7)	
Gender				0.741
Male	2185 (50.9)	1344 (50.6)	841 (51.6)	
Female	2070 (49.1)	1311 (49.4)	759 (48.4)	
BMI (kg/m ²)				0.113
< 25	528 (11.3)	311 (11.4)	217 (11.0)	
25–30	1191 (24.8)	787 (26.2)	404 (22.1)	
≥30	2536 (63.9)	1557 (62.4)	979 (66.9)	
Smoking				0.008
Never smoking	2140 (49.1)	1376 (50.8)	764 (45.8)	
Former smoking	1433 (35.1)	840 (32.2)	593 (40.5)	
Now smoking	682 (15.8)	439 (16.9)	243 (13.7)	
Alcohol drinking				0.008
No	1303 (26.5)	775 (24.5)	528 (30.3)	
Yes	2952 (73.5)	1880 (75.5)	1072 (69.7)	
Hypertension				<0.001
No	871 (21.0)	679 (25.8)	192 (11.9)	
Yes	3384 (79.0)	1976 (74.2)	1408 (88.1)	
Family history of diabetes				0.974
No	1510 (36.7)	964 (36.7)	546 (36.7)	
Yes	2745 (63.3)	1691 (63.3)	1054 (63.3)	
Physical activity				<0.001
None	902 (22.7)	624 (24.8)	278 (18.8)	
Moderate	1448 (36.3)	957 (38.3)	491 (32.6)	
Intense	1905 (41.0)	1074 (36.9)	831 (48.6)	
Waist (cm)	111.45 (16.31)	110.72 (16.29)	112.81 (16.28)	0.006
HDL (mg/dL)	47.27 (14.65)	47.80 (14.68)	46.26 (14.54)	0.018
TC (mg/dL)	186.26 (47.17)	186.85 (45.35)	185.13 (50.42)	0.367
BUN (mg/dL)	15.89 (7.27)	14.00 (4.61)	19.46 (9.65)	<0.001
Uric acid (mg/dL)	5.72 (1.46)	5.49 (1.28)	6.16 (1.67)	<0.001
HEI-2020	51.91 (11.60)	52.33 (11.72)	51.13 (11.32)	0.036

Note: DKD, diabetic kidney disease; BMI, body mass index; HDL, high-density lipoprotein cholesterol; TC, total cholesterol; BUN, blood urea nitrogen; HEI-2020, Healthy Eating Index-2020.
n (%) represented the categorical variable, and mean (SD) represented the continuous variable. n was unweighted, n (%), mean, and SD were weighted.

Table 2
Associations between HEI-2020 and odds ratios (95% confidence intervals) for DKD.

Participants	OR (95% CI), P-value		
	Crude	Model 1	Model 2
HEI-2020 (continuous)	0.991 (.983–0.999), 0.034	0.979 (0.970–0.988), <0.001	0.982 (0.973–0.992), <0.001
HEI-2020 (categorical)			
T1 (≤46.433)	Ref.	Ref.	Ref.
T2 (46.433–56.822)	0.928 (0.753–1.145), 0.484	0.751 (0.597–0.945), 0.015	0.767 (0.597–0.985), 0.038
T3 (> 56.822)	0.847 (0.677–1.060), 0.145	0.622 (0.488–0.793), <0.001	0.666 (0.518–0.857), 0.002
P for trend	0.145	<0.001	0.001

Note: HEI-2020, Healthy Eating Index-2020; OR, odds ratio; CI, confidence interval.
Crude model: without adjustment; Model 1: with adjustments on age, gender, and race; Model 2: with adjustments on age, gender, and race, BMI, smoking, drinking, physical activity, waist circumference, hypertension, family history of diabetes, high-density lipoprotein cholesterol, total cholesterol, blood urea nitrogen, uric acid.

in T2D patients. In a fully adjusted model, the HEI-2020 WQS index was significantly inversely correlated with the prevalence of DKD in T2D patients (OR = 0.765, 95% CI: 0.621–0.943, P = 0.012). Among these, dairy products (22.36%), total protein foods (22.10%), and total vegetables (10.99%) stood out as the most impactful components (Fig. 4).




Fig. 2. RCS diagram.

Characters	n (%)	OR (95%CI)	P value	P interaction
Gender				0.293
Male	2185 (50.9)	0.987 (0.973–1.000)	0.049	
Female	2070 (49.1)	0.976 (0.964–0.989)	<0.001	
BMI				0.877
<25	528 (11.3)	0.979 (0.953–1.005)	0.099	
25–30	1191 (24.8)	0.977 (0.960–0.994)	0.008	
≥30	2536 (63.9)	0.986 (0.973–0.998)	0.025	
Smoking				0.303
Never smoking	2140 (49.1)	0.977 (0.963–0.991)	0.001	
Former smoking	1433 (35.1)	0.981 (0.963–0.999)	0.034	
Now Smoking	682 (15.8)	1.004 (0.984–1.025)	0.692	
Alcohol drinking				0.875
No	1303 (26.5)	0.983 (0.968–0.999)	0.032	
Yes	2952 (73.5)	0.982 (0.970–0.993)	0.001	
Hypertension				0.895
No	871 (21.0)	0.982 (0.961–1.004)	0.099	
Yes	3384 (79.0)	0.982 (0.971–0.992)	0.001	
Physical activity				0.382
None	902 (22.7)	0.983 (0.961–1.005)	0.122	
Moderate	1448 (36.3)	0.988 (0.973–1.002)	0.097	
Intense	1905 (41.0)	0.977 (0.962–0.992)	0.003	
Family history of diabetes				0.628
No	1510 (36.7)	0.978 (0.963–0.993)	0.003	
Yes	2745 (63.3)	0.984 (0.972–0.996)	0.008	

Fig. 3. Subgroup forest plot.

Table 3
Independent effects of HEI-2020 related components on the prevalence of DKD in patients with type 2 diabetes.

Component	OR	95% CI	P-value
Total vegetables	0.916	0.841–0.999	0.043
Greens and beans	0.952	0.883–1.027	0.193
Total fruits	0.973	0.913–1.038	0.406
Whole fruits	0.977	0.924–1.032	0.395
Whole grains	0.994	0.961–1.029	0.739
Dairy	0.953	0.911–0.997	0.032
Total protein foods	0.884	0.796–0.981	0.018
Seafood and plant proteins	0.902	0.839–0.970	0.005
Fatty acids	0.985	0.945–1.026	0.461
Sodium	0.999	0.957–1.043	0.968
Refined grains	0.965	0.930–1.001	0.053
Saturated fats	0.992	0.945–1.042	0.747
Added sugars	0.937	0.888–0.989	0.017

Note: OR, odds ratio; CI, confidence interval.

Relationship between HEI-2020 and DKD in newly diagnosed and previously diagnosed T2D population

Considering the important impact of the duration of diabetes on DKD,²⁰ we investigated the relationship between HEI-2020 and DKD in newly diagnosed and previously diagnosed people with T2D. Among 4255 participants, 3125 patients with T2D were previously diagnosed, and the prevalence of DKD was 40.9%. There were 1130 newly diagnosed T2D patients, and the prevalence of DKD was 28.6%. The results of the weighted logistic regression model showed that, after adjusting all covariates, HEI-2020 scores were significantly negatively correlated with DKD in previously diagnosed T2D patients (OR = 0.979, 95% CI: 0.967–0.990, $P < 0.001$). Among them, the total protein food score (OR = 0.870, 95% CI: 0.769–0.984, $P = 0.024$), seafood and plant protein score (OR = 0.874, 95% CI:




Fig. 4. Estimated weight of WQS index.

0.806–0.948, $P = 0.001$), and refined grain score ($OR = 0.945$, 95% $CI: 0.904–0.987$, $P = 0.010$) were all associated with a lower risk of DKD. However, no significant association between HEI-2020 and DKD was observed in newly diagnosed T2D patients (Table 4).

Furthermore, subgroup analysis was carried out in previously diagnosed T2D patients. We found that the negative correlation

Table 4
Association between HEI-2020 and DKD in newly diagnosed and previously diagnosed type 2 diabetes populations.

Participants	OR (95% CI), P-value Model 2
Prior diagnosis	
HEI-2020	0.979 (0.967–0.990), <0.001
HEI-2020 component	
Total vegetables	0.930 (0.845–1.023), 0.130
Greens and beans	0.941 (0.866–1.022), 0.144
Total fruits	0.978 (0.903–1.059), 0.573
Whole fruits	0.966 (0.900–1.036), 0.321
Whole grains	0.967 (0.924–1.011), 0.135
Dairy	0.969 (0.922–1.019), 0.217
Total protein foods	0.870 (0.769–0.984), 0.024
Seafood and plant proteins	0.874 (0.806–0.948), 0.001
Fatty acids	0.968 (0.925–1.013), 0.154
Sodium	0.975 (0.924–1.029), 0.357
Refined grains	0.945 (0.904–0.987), 0.010
Saturated fats	0.992 (0.939–1.047), 0.759
Added sugars	0.973 (0.908–1.042), 0.422
New diagnosis	
HEI-2020	0.992 (0.975–1.010), 0.364
HEI-2020 component	
Total vegetables	0.877 (0.754–1.020), 0.083
Greens and beans	0.960 (0.833–1.108), 0.572
Total fruits	0.969 (0.861–1.090), 0.594
Whole fruits	1.016 (0.909–1.135), 0.780
Whole grains	1.081 (0.998–1.170), 0.051
Dairy	0.900 (0.818–0.991), 0.029
Total protein foods	0.880 (0.707–1.095), 0.244
Seafood and plant proteins	0.940 (0.818–1.081), 0.380
Fatty acids	1.028 (0.949–1.114), 0.488
Sodium	1.076 (0.998–1.160), 0.054
Refined grains	1.029 (0.951–1.114), 0.467
Saturated fats	0.995 (0.913–1.085), 0.913
Added sugars	0.846 (0.789–0.909), <0.001

Model 2: with adjustments on age, gender, and race, BMI, smoking, drinking, physical activity, waist circumference, hypertension, family history of diabetes, high-density lipoprotein cholesterol, total cholesterol, blood urea nitrogen, uric acid. Note: HEI-2020, Healthy Eating Index-2020; OR, odds ratio; CI, confidence interval.

between HEI-2020 score and DKD was significant in most subgroups. The interaction test showed that there was a significant interaction between smoking status (P for interaction = 0.037). A higher HEI-2020 was significantly associated with a lower risk of DKD in individuals who never smoked or had previously smoked. However, no statistical association was observed among the population who were current smokers (Table S2).

Sensitivity analysis

On the basis of Model 2, we further adjusted ACEIs, ARBs, SGLT2 inhibitors, GLP-1 receptor agonists, and statins. The results showed that HEI-2020 was still significantly negatively correlated with the risk of DKD in patients with T2D ($OR = 0.982$, 95% $CI: 0.973–0.991$, $P < 0.001$). The scores of total vegetables, dairy products, total protein foods, seafood and plant protein, refined grains, and added sugars were significantly negatively correlated with the incidence of DKD (Table S3), which was highly consistent with our main analysis and demonstrated the robustness of our results.

Discussion

Our findings demonstrated a significant negative correlation between HEI-2020 scores and the prevalence of DKD among T2D patients. This association was observed in individuals who were overweight, obese, non-smoking, had previously smoked, had high blood pressure, and engaged in intense PA. Furthermore, dairy products, total protein foods, and total vegetables were the most impactful components in HEI-2020.

HEI has been considered to be associated with the prevalence of diabetes and CKD. Specifically, the HEI-2005 score is positively correlated with reduced risk of diabetes,⁹ while the HEI-2010 score was discovered to be negatively correlated with microalbuminuria concentration in diabetic patients.²¹ A prospective cohort study revealed that compared with those in the lowest quintile, participants in the highest quintile of the HEI-2015 score had a 17% lower risk of developing CKD,²² and the highest quintile was also associated with a composite risk of kidney outcomes reduction.²³ The new version of the HEI-2020 showed a significant negative correlation with the prevalence of DKD in T2D patients, further emphasizing the favorable effect of HEI scoring. The components and scorings of different versions of HEI are shown in Table S4. Furthermore, HEI-2020 is associated with the risk of heart failure and reduced levels of inflammatory markers,²⁴ which exacerbate DKD progression.²⁵ Therefore, for individuals with diabetes, adhering to the healthy eating patterns advocated by HEI-2020 not only aids in lowering the risk of DKD but may also positively impact the prevention and control of other chronic diseases, thereby improving overall well-being.

In different populations characterized by distinct features, the HEI-2020 score exhibited disparities in the prevalence of DKD among patients with T2D. While metabolic abnormalities may be more pronounced in overweight and obese patients, the prevalence of DKD is potentially higher.²⁶ However, increased HEI scores can alleviate weight gain and enhance overall health.²⁷ Therefore, it may be more beneficial for such patients. For those with hypertension, HEI emphasizes limiting sodium intake and consuming more vegetables and fruits, which effectively reduces the risk of hypertension,²⁸ exhibiting potential renal protective effects. High-intensity exercise can improve blood sugar control, reduce renal oxidative stress and inflammation, and synergistically improve risk factors such as obesity and hypertension with HEI diet, thereby being associated with a lower risk of DKD.²⁹ In addition, high-intensity interval training is superior to continuous training with moderate intensity in controlling blood glucose control.³⁰ On the contrary, patients who lack exercise often have more metabolic abnormalities, obesity, and sedentary behavior,

which increases their risk of DKD,³¹ and possibly masks the true association between HEI-2020 and DKD risk.

Upon further analysis, it was found that dairy products, total protein foods, and total vegetables were the most influential components within the HEI-2020. Dairy products are a class of food made from milk, which have numerous health benefits for the human body.³² Research shows that higher intake of milk and yogurt is associated with lower risk of DKD and metabolic syndrome.³³ Probiotic yogurt (such as kefir) can also lower blood glucose, creatinine, and urea levels in diabetic rats, and improve renal glomerular casts, tubular degeneration, and renal epithelial tissue, thereby slowing down the progression of DKD.³⁴ In addition, camel milk has been shown to downregulate the expression of DKD-related genes and preserve normal glomerular morphology.³⁵

Protein is an indispensable nutrient for human life activities and physiological functions. A higher protein intake is associated with a lower prevalence of prediabetes and diabetes,³⁶ as well as improved blood lipid levels and insulin resistance in T2D.³⁷ Protein intake can augment satiety, reduce lipid absorption in the gut, and repress fat deposition by curbing the growth of *Lactobacilli*,³⁸ thereby aiding in appetite control and weight maintenance. These changes are all associated with a decrease in the incidence of DKD.

The consumption of total vegetables also holds significant sway over the prevalence of DKD. Genetic evidence suggests that vegetable intake is determined as a protective factor for DKD.³⁹ A case-control study also demonstrated that vegetable consumption was associated with lower DKD prevalence.⁴⁰ Dietary fiber can regulate gut microbiota, enrich bacteria that secrete short-chain fatty acids (such as *Prevotella* and *Bifidobacterium*), and increase short-chain fatty acids. It can also reduce the expression of genes encoding inflammatory cytokines and chemokines in DKD, thereby preventing the occurrence of DKD.⁴¹ In addition, the antioxidants found in vegetables, including vitamin E, carotenoids, and vitamin C, can effectively alleviate oxidative stress and chronic inflammation associated with diabetes,⁴² thereby contributing to slowing down the development of DKD.

We also found that consuming more seafood and plant protein was associated with a lower risk of DKD, while consuming added sugar was associated with a higher risk of DKD. Seafood products are rich in omega-3 fatty acids, which have anti-inflammatory and antioxidant properties.⁴³ Increased intake of omega-3 fatty acids can lower blood pressure, which is a key risk factor for the development of CKD.⁴⁴ Regarding plant protein, research has shown that plant or white meat protein is more beneficial for kidney function protection compared to red meat protein.⁴⁵ Beans are an important source of plant protein. A study compared the effects of different soy protein intake ratios (0%, 35%, 100%) on DKD patients, and the results showed that a 35% soy protein diet was most effective in improving kidney function and glucose and lipid metabolism.⁴⁶ Added sugar is commonly believed to cause insulin resistance, increase oxidative stress and inflammation levels, thereby impairing kidney function, which is consistent with our findings.⁴⁷ Therefore, it is advisable to limit the intake of added sugars as much as possible, which can help maintain kidney health.

This study further explored the impact of diabetes duration on the association between HEI-2020 and DKD. In the previously diagnosed T2D patients, HEI-2020 was significantly negatively correlated with DKD, while no significant correlation was observed in the newly diagnosed patients, which suggested that the course of diabetes may affect the effect of diet-related interventions. The longer duration of diabetes is proven to be associated with more obvious renal damage caused by chronic hyperglycemia.²⁰ Long-term adherence to a healthy diet may alleviate the sustained adverse effects of high blood sugar on the kidneys to some extent by improving metabolic status, reducing inflammation levels, and oxidative stress.⁴⁸ In contrast, the risk of DKD in newly diagnosed patients is mainly influenced by genetic factors and other comorbid risk factors, and the effect of dietary intervention may not have been fully demonstrated. Subgroup

analysis of previously diagnosed T2D patients found that smoking status had a significant interaction on the association between HEI-2020 and DKD. Smoking is a known risk factor for DKD, and nicotine in it will increase the expression of proteinuria and fibronectin in diabetic mice.⁴⁹ This suggests that current smoking may weaken the protective effect of a healthy diet on the kidneys, thereby offsetting the potential benefits of a healthy diet. It is worth noting, however, that some limitations in the research should also come into consideration. Since this work is a cross-sectional analysis, we can only identify the relationship between variables rather than establish causality. Secondly, according to the KDIGO guidelines, CKD is defined as persistent abnormalities in renal structure or function (eGFR < 60 mL/min/1.73 m² and/or urinary albumin/creatinine ratio ≥ 30 mg/g) that persist for ≥ 3 months.⁵⁰ In this study, we only evaluated renal function based on a single measurement of eGFR and urinary albumin/creatinine ratio. The transient renal dysfunction may increase the risk of overestimating the prevalence of DKD. Thirdly, this study excluded a large number of samples due to missing covariate information from some participants, which may lead to selection bias and affect the representativeness of the results. Fourthly, all participants are drawn from the NHANES program, which may limit the generalizability of our findings when applied to other populations. Fifthly, despite our efforts to adjust potential confounders in the analysis process to minimize their impact on the results, we cannot entirely exclude the impact of other potential confounders such as environmental factors (air quality and water quality), metabolic levels, as well as genetic structures. In the present study, our main focus was on the impact of HEI-2020 scores on the prevalence of DKD. Unfortunately, we did not delve into an in-depth analysis regarding whether the long-term prognosis of patients with DKD would change. Future studies can dig out the impact of HEI-2020 scores on the outcomes of DKD patients.

Conclusion

This study revealed a negative correlation between HEI-2020 scores and DKD prevalence among T2D patients. Additionally, dairy products, complete protein foods, and total vegetables offer significant health benefits in reducing the prevalence of DKD. These findings underscore the importance of dietary management in T2D patients, suggesting that personalized dietary strategies could facilitate the health of such individuals and mitigate the risk of DKD.

CRediT authorship contribution statement

Conceptualization: Chunping Dong, Hui Li, Jing Li; Data curation: Hui Li; Formal analysis: Chunping Dong, Shan Gao; Investigation: Jing Li; Methodology: Chunping Dong; Project administration: Chunping Dong; Resources: Hui Li; Software: Yuan Qiao; Supervision: Hui Li; Validation: Shan Gao; Visualization: Shan Gao; Writing – original draft: Chunping Dong, Hui Li, Jing Li; Writing – review & editing: Yuan Qiao, Shan Gao.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Funding

Not applicable.

Conflict of interest

Not applicable.

Data availability

The data and materials in the current study are available from the corresponding author on reasonable request.

Appendix A. Supplementary data

Supplementary data associated with this article can be found in the online version available at <https://doi.org/10.1016/j.nefro.2025.501452>.

References

- Kolaric V, Svircevic V, Bijuk R, Zupancic V. Chronic complications of diabetes and quality of life. *Acta Clin Croat*. 2022;61:520–7.
- IDF The Diabetes Atlas. Available from: <https://diabetesatlas.org/>.
- van Raalte DH, Bjornstad P, Cherney DZ, de Boer I, Fioretto IH, Gordin P, et al. Combination therapy for kidney disease in people with diabetes mellitus. *Nat Rev Nephrol*. 2024;20:433–46.
- Wen-Yue L, Wen-Ying C, Jia-Hui Z, Giovanni T, Christopher DB, Anoop M, et al. The global burden of diabetes-related chronic kidney disease from 1990 to 2021, with projections to 2036. *Metab Target Organ Damage*. 2025;5:41.
- Gonzalez-Perez A, Saez M, Vizcaya D, Lind M, Garcia Rodriguez L. Incidence and risk factors for mortality and end-stage renal disease in people with type 2 diabetes and diabetic kidney disease: a population-based cohort study in the UK. *BMJ Open Diabetes Res Care*. 2021;9.
- Gupta S, Dominguez M, Golestaneh L. Diabetic kidney disease: an update. *Med Clin North Am*. 2023;107:689–705.
- Real Rodriguez CC, Riboldi BP, Rodrigues TD, Sarmiento C, Antonio RA, de Almeida JPJC. Association of eating patterns and diabetic kidney disease in type 2 diabetes: a cross-sectional study. *J Ren Nutr*. 2023;33:261–8.
- Shams-White MM, Pannucci TE, Lerman JL, Herrick KA, Zimmer M, Meyers Mathieu K, et al. Healthy Eating Index-2020: review and update process to reflect the Dietary Guidelines for Americans, 2020–2025. *J Acad Nutr Diet*. 2023;123:1280–8.
- Chiuvè SE, Fung TT, Rimm EB, Hu FB, McCullough ML, Wang M, et al. Alternative dietary indices both strongly predict risk of chronic disease. *J Nutr*. 2012;142:1009–18.
- Brlek A, Gregoric M. Diet quality indices and their associations with all-cause mortality CVD and type 2 diabetes mellitus: an umbrella review. *Br J Nutr*. 2023;130:709–18.
- Zeinalabedini M, Nasli-Esfahani E, Esmailzadeh A, Azadbakht L. How is healthy eating index-2015 related to risk factors for cardiovascular disease in patients with type 2 diabetes. *Front Nutr*. 2023;10. 1201010.
- Huang Y, Xu S, Wan T, Wang X, Jiang S, Shi W, et al. The combined effects of the most important dietary patterns on the incidence and prevalence of chronic renal failure: results from the US National Health and Nutrition Examination Survey and Mendelian analyses. *Nutrients*. 2024;16.
- American Diabetes Association. 2. Classification and diagnosis of diabetes: standards of medical care in diabetes – 2019. *Diabetes Care*. 2019;42 Suppl. 1:S13–28.
- Caspard H, Jabbour S, Hammar N, Fenici P, Sheehan JJ, Kosiborod M. Recent trends in the prevalence of type 2 diabetes and the association with abdominal obesity lead to growing health disparities in the USA: an analysis of the NHANES surveys from 1999 to 2014. *Diabetes Obes Metab*. 2018;20:667–71.
- Krebs-Smith SM, Pannucci TE, Subar AF, Kirkpatrick SL, Lerman JL, Toozé JA, et al. Update of the Healthy Eating Index: HEI-2015. *J Acad Nutr Diet*. 2018;118:1591–602.
- Kidney Disease: Improving Global Outcomes Glomerular Diseases Work Group. KDIGO 2021 clinical practice guideline for the management of glomerular diseases. *Kidney Int*. 2021;100:S1–276.
- Zhou C, Zhou Y, Shuai N, Zhou J, Kuang X. The nonlinear relationship between estimated glomerular filtration rate and cardiovascular disease in US adults: a cross-sectional study from NHANES 2007–2018. *Front Cardiovasc Med*. 2024;11. 1417926.
- Chen H, Leng X, Liu S, Zeng Z, Huang F, Huang R, et al. Association between dietary intake of omega-3 polyunsaturated fatty acids and all-cause and cardiovascular mortality among hypertensive adults: results from NHANES 1999–2018. *Clin Nutr*. 2023;42:2434–42.
- Neuen BL, Fletcher RA, Heath L, Perkovic A, Vaduganathan M, Badve SV, et al. Cardiovascular kidney, and safety outcomes with GLP-1 receptor agonists alone and in combination with SGLT2 inhibitors in type 2 diabetes: a systematic review and meta-analysis. *Circulation*. 2024;150:1781–90.
- Siddiqui K, George TP, Joy SS, Alfadda AA. Risk factors of chronic kidney disease among type 2 diabetic patients with longer duration of diabetes. *Front Endocrinol (Lausanne)*. 2022;13. 1079725.
- Costacou T, Crandell J, Kahkoska AR, Liese AD, Dabelea D, Lawrence JM, et al. Dietary patterns over time and microalbuminuria in youth and young adults with type 1 diabetes: the SEARCH Nutrition Ancillary Study. *Diabetes Care*. 2018;41:1615–22.
- Hu EA, Steffen LM, Grams ME, Crews DC, Coresh J, Appel LJ, et al. Dietary patterns and risk of incident chronic kidney disease: the atherosclerosis risk in communities study. *Am J Clin Nutr*. 2019;110:713–21.
- Smyth A, Griffin M, Yusuf S, Mann JF, Reddan D, Canavan M, et al. Diet and major renal outcomes: a prospective cohort study. The NIH-AARP Diet and Health Study. *J Ren Nutr*. 2016;26:288–98.
- Gu F, Yu W, Shu T, Zhu Y. Association between the healthy eating index 2020 and heart failure among the U.S. middle-aged and older adults from NHANES 2005–2020: a cross-sectional study. *Front Nutr*. 2024;11. 1496379.
- Guo W, Song Y, Sun Y, Du H, Cai Y, You Q, et al. Systemic immune-inflammation index is associated with diabetic kidney disease in type 2 diabetes mellitus patients: evidence from NHANES 2011–2018. *Front Endocrinol (Lausanne)*. 2022;13. 1071465.
- Zhang K, Zhang W, Xia F, Wang N, Lu Y, Sui C, et al. Obesity patterns metabolic abnormality, and diabetic kidney disease in patients with type 2 diabetes. *Diabetes Metab Syndr Obes*. 2023;16:3999–4011.
- Vahid F, Jalili M, Rahmani W, Nasiri Z, Bohn T. A higher healthy eating index is associated with decreased markers of inflammation and lower odds for being overweight/obese based on a case-control study. *Nutrients*. 2022;14.
- Pasdar Y, Hamzeh B, Moradi S, Mohammadi E, Cheshmeh S, Darbandi M, et al. Healthy eating index 2015 and major dietary patterns in relation to incident hypertension: a prospective cohort study. *BMC Public Health*. 2022;22:734.
- Fan R, Kong J, Zhang J, Zhu L. Exercise as a therapeutic approach to alleviate diabetic kidney disease: mechanisms, clinical evidence and potential exercise prescriptions. *Front Med (Lausanne)*. 2024;11. 1471642.
- Mendes R, Sousa N, Themudo-Barata JL, Reis VM. High-intensity interval training versus moderate-intensity continuous training in middle-aged and older patients with type 2 diabetes: a randomized controlled crossover trial of the acute effects of treadmill walking on glycemic control. *Int J Environ Res Public Health*. 2019;16.
- Kerr NR, Booth FW. Contributions of physical inactivity and sedentary behavior to metabolic and endocrine diseases. *Trends Endocrinol Metab*. 2022;33:817–27.
- Tunick MH, Van Hekken DL. Dairy products and health: recent insights. *J Agric Food Chem*. 2015;63:9381–8.
- Yoshinari M, Ohkuma T, Iwase M, Kitazono T. Milk and yogurt consumption and its association with cardiometabolic risk factors in patients with type 2 diabetes: The Fukuoka Diabetes Registry. *Nutr Metab Cardiovasc Dis*. 2025;35. 103772.
- Kahraman M, Ertekin YH, Satman I. The effects of kefir on kidney tissues and functions in diabetic rats. *Probiotics Antimicrob Proteins*. 2021;13:375–82.
- Shaban AM, Raslan M, Qahl SH, Elsayed K, Abdelhameed MS, Oyouni AA, et al. Ameliorative effects of camel milk and its exosomes on diabetic nephropathy in rats. *Membranes (Basel)*. 2022;12.
- Sluik D, Brouwer-Brolsma EM, Berendsen AA, Mikkila M, Poppitt V, Silvestre SD, et al. Protein intake and the incidence of pre-diabetes and diabetes in 4 population-based studies: the PREVIEW project. *Am J Clin Nutr*. 2019;109:1310–8.
- Yu Z, Nan F, Wang LY, Jiang H, Chen W, Jiang Y. Effects of high-protein diet on glycemic control, insulin resistance and blood pressure in type 2 diabetes: a systematic review and meta-analysis of randomized controlled trials. *Clin Nutr*. 2020;39:1724–34.
- Zhong W, Wang H, Yang Y, Zhang Y, Lai H, Cheng Y, et al. High-protein diet prevents fat mass increase after dieting by counteracting *Lactobacillus*-enhanced lipid absorption. *Nat Metab*. 2022;4:1713–31.
- Zhou Y, Liu Y, Wu L, Zhang Y, Wen H, Hu J, et al. Causal insights into major risk factors for diabetic kidney disease: a comprehensive meta-analysis and Mendelian randomization study. *Ren Fail*. 2025;47. 2468741.
- Rezazadegan M, Mirjalili F, Jalilpiran Y, Aziz M, Jayedi A, Setayesh L, et al. The association between dietary diversity score and odds of diabetic nephropathy: a case-control study. *Front Nutr*. 2022;9. 767415.
- Li YJ, Chen X, Kwan TK, Loh YW, Singer J, Liu Y, et al. Dietary fiber protects against diabetic nephropathy through short-chain fatty acid-mediated activation of G protein-coupled receptors GPR43 and GPR109A. *J Am Soc Nephrol*. 2020;31:1267–81.
- Gowd V, Xiao J, Wang M, Chen F, Cheng KW. Multi-mechanistic antidiabetic potential of astaxanthin: an update on preclinical and clinical evidence. *Mol Nutr Food Res*. 2021;65. e2100252.
- Patted PG, Masareddy RS, Patil AS, Kanabargi RR, Bhat CT. Omega-3 fatty acids: a comprehensive scientific review of their sources, functions and health benefits. *Future J Pharm Sci*. 2024;10:94.
- Zhang X, Ritonja JA, Zhou N, Chen BE, Li X. Omega-3 polyunsaturated fatty acids intake and blood pressure: a dose-response meta-analysis of randomized controlled trials. *J Am Heart Assoc*. 2022;11. e025071.
- Eckert I, Koehler IC, Bauer J, Busnello FM, Silva FM. Effects of different sources of dietary protein on markers of kidney function in individuals with diabetes: a systematic review and meta-analysis of randomized controlled trials. *Nutr Rev*. 2022;80:812–25.

46. Sun J, Wei Y, Miao R, Zhang X, Zhang B, Zhang L, et al. Comparison of the effects of different percentages of soy protein in the diet on patients with type 2 diabetic nephropathy: systematic reviews and network meta-analysis. *Front Nutr.* 2023;10:1184337.

47. DiNicolantonio JJ, O’Keefe JH, Bhutani J. Added sugars drive chronic kidney disease and its consequences: a comprehensive review. *J Metab Health.* 2016;1:1–6.

48. Hu H, Ding G, Liang W. Dietary therapy to halt the progression of diabetes to diabetic kidney disease. *Food Funct.* 2025;16:2622–36.

49. Jaimes EA, Zhou MS, Siddiqui M, Rezonzew G, Tian R, Seshan SV, et al. Nicotine, smoking, podocytes, and diabetic nephropathy. *Am J Physiol Renal Physiol.* 2021;320:F442–53.

50. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2024 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int.* 2024;105:S117–314.

1059
1060
1061
1062
1063
1064

UNCORRECTED PROOF