

BIBLIOGRAFÍA

- Bilgin H, Sarmis A, Tigen E, Soyletir G, Mulazimoglu L. Delftia acidovorans: A rare pathogen in immunocompetent and immunocompromised patients. *Can J Infect Dis Med Microbiol.* 2015;26:277-9.
- Mahmood S, Taylor KE, Overman TL, McCormick MI. Acute infective endocarditis caused by Delftia acidovorans, a rare pathogen complicating intravenous drug use. *J Clin Microbiol.* 2012;50:3799-800.
- Horowitz H, Gilroy S, Feinstein S, Gilardi G. Endocarditis associated with Comamonas acidovorans. *J Clin Microbiol.* 1990;28:143-5.
- Yildiz H, Sünnetçioğlu A, Ekin S, Baran Aİ, Özgökçe M, Aşker S, et al. Delftia acidovorans pneumonia with lung cavities formation. *Colomb Med (Cali).* 2019;50:215-21.
- Khan S, Sistla S, Dhodapkar R, Parija SC. Fatal Delftia acidovorans infection in an immunocompetent patient with empyema. *Asian Pac J Trop Biomed.* 2012;2:923.
- Deb AK, Chavhan P, Chowdhury SS, Sistla S, Sugumaran S, Panicker G. Endophthalmitis due to Delftia acidovorans: An unusual ocular pathogen. *Indian J Ophthalmol.* 2020;68:2591-4.
- Sahay P, Goel S, Nagpal R, Maharana PK, Sinha R, Agarwal T, et al. Infectious Keratitis Caused by Rare and Emerging Micro-Organisms. *Curr Eye Res.* 2020;45:761-73.
- Lee SM, Kim MK, Lee JL, Wee WR, Lee JH. Experience of Comamonas acidovorans keratitis with delayed onset and treatment response in immunocompromised cornea. *Korean J Ophthalmol.* 2008;22:49-52.
- López-Menchero R, Sigüenza F, Caridad A, Alonso JC, Ferreruela RM. Peritonitis due to Comamonas acidovorans in a CAPD patient. *Perit Dial Int.* 1998;18:445-56.
- Calzada M, Roig M, Martínez-Toldos MC, Segovia M. Urinary tract infection associated with Delftia acidovorans. *Rev Esp Quimioter.* 2015;28:326-7.
- Ojeda-Vargas MM, Suárez-Alonso A, Pérez Cervantes MA, Suárez-Gil E, Monzón-Moreno C. Urinary tract infection associated with Comamonas acidovorans. *Clin Microbiol Infect.* 1999;5:443-4.
- Chotikanatis K, Martin Bäcker M, Gabriela Rosas-Garcia G, Hammerschlag MR. Recurrent intravascular-catheter-related bacteremia caused by Delftia acidovorans in a hemodialysis patient. *J Clin Microbiol.* 2011;49:3418-21.
- Kawamura I, Yagi T, Hatakeyama K, Ohkura T, Ohkusu K, Takahashi Y, et al. Recurrent vascular catheter-related bacteremia caused by Delftia acidovorans with different antimicrobial susceptibility profiles. *J Infect Chemother.* 2010;17:111-3.
- Lipuma JJ, Currie BJ, Peacock SJ, Vandamme P, Whittier S. Burkholderia, Cupriavidus, Pandoraea, Stenotrophomonas, Ralstonia, Brevundimonas, Comamonas Delftia, and Acidovorax. *Manual of Clinical Microbiology.* 10th edn. Washington, DC: ASM Press; 2011. p. 692-713.
- Kam SK, Lee WS, Ou TS, Teng SO, Chen FL. Delftia acidovorans Bacteremia Associated with Ascending Urinary Tract Infections Proved by Molecular Method. *J Exp Clin Med.* 2012;4:180-2.
- Højgaard SMM, Reza Hosseini O, Knudsen JD, Fuglebjerg NJU, Skov M, Nielsen SD, et al. Characteristics and Outcomes of Patients with Delftia acidovorans Infections: a Retrospective Cohort Study. *Microbiol Spectr.* 2022;10:e0032622.

Noa Díaz Novo^{a,*}, Daniel Adrados Ruiz^b,
Beatriz Crespo Estrada^b, Ingrid Auyanet Saavedra^a,
Ana Ramírez Puga^a, Rita Guerra Rodríguez^a,
Ernesto Fernández-Tagarro^a
y César García-Cantón^a

^a Servicio de Nefrología, Hospital Universitario Insular de Gran Canaria, Las Palmas de Gran Canaria, España

^b Servicio de Microbiología, Hospital Universitario Insular de Gran Canaria, Las Palmas de Gran Canaria, España

* Autor para correspondencia.

Correo electrónico: ndianov@gobiernodecanarias.org (N. Díaz Novo).

0211-6995/© 2023 Sociedad Española de Nefrología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.nefro.2022.12.010>

Targeting colonic BK channels: A novel therapeutic strategy against hyperkalemia in chronic kidney disease

Canales BK colónicos: una nueva estrategia terapéutica contra la hiperpotasemia en la enfermedad renal crónica



Dear Editor,

Hyperkalemia is a common electrolyte disorder frequently observed in patients with chronic kidney disease (CKD).¹ Once potassium (K⁺) ions are absorbed from the daily food intake, they are predominantly distributed in the intracellular space

than in the extracellular space.¹ Under physiological conditions, approximately 90% of the dietary K⁺ ions are excreted into the urine, while the remaining 10% are excreted into the feces. The prompt management of hyperkalemia is to temporarily shift the extracellular K⁺ ions into the intracellular




Fig. 1 – BK channels in colonic enterocytes as the therapeutic target for hyperkalemia in chronic kidney disease (CKD). BK channels are expressed in the apical membrane of colonic enterocytes and play a major role in excreting K^+ ions into the feces. Already marketed drugs, such as anti-platelets (cilostazol) and anti-epileptics (zonisamide), or natural compounds, such as estradiol, omega-3 docosahexaenoic acid (DHA) and resveratrol, increase the BK channel activity. Osmotic laxatives, which increase the intestinal luminal flow, also activate the BK channels and thus ameliorate hyperkalemia in CKD.

space by insulin.¹ However, to fundamentally treat hyperkalemia, the redundant K^+ ions have to be eliminated from the body, either by the use of diuretics to facilitate the renal K^+ excretion or by performing emergent hemodialysis.¹

In the kidneys, two populations of K^+ channels are expressed in the apical membrane of the collecting duct principal cells.² These channels, such as the renal outer medullary K^+ channels (ROMK or $Kir_{1.1}$) and large-conductance calcium-activated potassium channels (BK or $Kca_{1.1}$), play major roles in excreting K^+ ions into the urine. Both K^+ channels are directly stimulated by aldosterone and also by the decrease in the luminal K^+ concentration and the increase in the distal sodium (Na^+) delivery.² Of note, BK channels are additionally stimulated by the increased renal tubular flow,³ since this elevates the intracellular calcium (Ca^{2+}) concentration, which is the primary trigger of the channel activation. In patients with CKD, due to the renal tubular damage or dysfunction,¹ these K^+ channels are frequently disabled. Consequently, the impaired renal K^+ excretion leads to the much higher prevalence of hyperkalemia than in the general population.¹

Similar to the renal collecting ducts, BK channels are also expressed in the apical membrane of colonic enterocytes and play a major role in excreting K^+ ions into the feces⁴ (Fig. 1). In diseases that cause secretory diarrhea, such as ulcerative colitis, enteric infections, villous adenoma, BK channels are over-expressed or over-activated in the colonic epithelial cells,⁴ resulting in the increased fecal K^+ losses. In CKD, to compensate for the decreased renal K^+ excretion, its fecal excretion was alternatively increased.⁵ In clinical stu-

dies, since colonic BK channels were actually over-expressed in patients with advanced CKD or end-stage renal disease (ESRD),⁵ these channels were thought to be primarily responsible for the increased fecal K^+ excretion. Therefore, drugs that directly stimulate the colonic BK channels would be effective in the treatment of hyperkalemia in CKD.

So far, several BK channel openers, including NS1619, NS11021 and BMS204352, have newly been synthesized.⁶ Additionally, recent studies revealed that already marketed drugs, such as anti-platelets (cilostazol) and anti-epileptics (zonisamide)⁶ or natural compounds, such as estradiol, omega-3 docosahexaenoic acid (DHA) and resveratrol,^{6,7} also increase BK channel activity (Fig. 1). Besides the kidney and distal intestine, BK channels are also expressed in the cardiovascular system and central nervous system.⁶ Therefore, in using BK channel openers, the adverse effects on these organs need to be carefully monitored. However, compared to the highly selective, newly synthesized BK channel openers, the already marketed drugs or natural compounds could be used more harmlessly, because they have commonly been used in clinical practices or consumed in daily diets for longer periods of time. To increase the efficacy of these drugs or compounds after orally administered, they need to target the colonic BK channels as specifically as possible. In this regard, recently developed “drug delivery systems” that effectively reach the colon may be beneficial.⁸ They include pH sensitive system, microbially triggered system, prodrugs, polysaccharide-based system, timed-release system, osmotically controlled drug system and pressure-dependent release system.⁸

In CKD patients, chronic constipation is one of the risk factors for developing hyperkalemia,⁹ since it mechanically disturbs the fecal K⁺ excretion. However, dissolving the constipation by probiotics alone did not ameliorate hyperkalemia in these patients. One of the commonly used laxatives, bisacodyl, is also a stimulator of BK channels,¹⁰ since it pharmacologically activates a cyclic adenosine monophosphate (cAMP). In CKD patients with hyperkalemia, the treatment with bisacodyl not only improved constipation, but also increased fecal K⁺ excretion and actually ameliorated hyperkalemia.¹⁰ Among laxatives, osmotic laxatives, such as magnesium hydroxide, polyethylene glycol, sorbitol and lactulose, exert their properties by drawing water into the colon and allowing easier and faster passage of the stool. In addition to the passive paracellular K⁺ excretion,¹⁰ such-induced increase in the intestinal luminal flow would activate the colonic BK channels (Fig. 1), similarly to the increased renal tubular flow that activates the renal BK channels.³ Given such pharmacological properties, the osmotic laxatives, which increase fecal K⁺ excretion through the activation of colonic BK channels, would also be beneficial in the treatment of hyperkalemia in CKD patients.

Conflict of interest

None declared.

Acknowledgements

This work was supported by the Salt Science Research Foundation, No. 2218 to IK.

BIBLIOGRAFÍA

1. Clase CM, Carrero JJ, Ellison DH, Grams ME, Hemmelgarn BR, Jardine MJ, et al. Potassium homeostasis and management of

dyskalemia in kidney diseases: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) controversies conference. *Kidney Int.* 2020;97:42–61.

2. Palmer BF. Regulation of potassium homeostasis. *Clin J Am Soc Nephrol.* 2015;10:1050–60.
3. Woda CB, Bragin A, Kleyman TR, Satlin LM. Flow-dependent K⁺ secretion in the cortical collecting duct is mediated by a maxi-K channel. *Am J Physiol Renal Physiol.* 2001;280:F786–93.
4. Rajendran VM, Sandle GI. Colonic potassium absorption and secretion in health and disease. *Compr Physiol.* 2018;8:1513–36.
5. Mathialahan T, MacLennan KA, Sandle LN, Verbeke C, Sandle GI. Enhanced large intestinal potassium permeability in end-stage renal disease. *J Pathol.* 2005;206:46–51.
6. Bentzen BH, Olesen SP, Ronn LC, Grunnet M. BK channel activators and their therapeutic perspectives. *Front Physiol.* 2014;5:389.
7. Li HF, Chen SA, Wu SN. Evidence for the stimulatory effect of resveratrol on Ca²⁺-activated K⁺ current in vascular endothelial cells. *Cardiovasc Res.* 2000;45:1035–45.
8. Bansal V, Malviya R, Malaviya T, Sharma PK. Novel prospective in colon specific drug delivery system. *Polim Med.* 2014;44:109–18.
9. Sumida K, Yamagata K, Kovesdy CP. Constipation in CKD. *Kidney Int Rep.* 2020;5:121–34.
10. Mathialahan T, Sandle GI. Dietary potassium and laxatives as regulators of colonic potassium secretion in end-stage renal disease. *Nephrol Dial Transplant.* 2003;18:341–7.

Itsuro Kazama*, Michiko Shoji

Miyagi University, School of Nursing, Gakuen, Taiwa-cho, Kurokawa-gun, Miyagi, Japan

* Corresponding author.

E-mail address: kazamai@myu.ac.jp (I. Kazama).

0211-6995/© 2023 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.nefro.2023.01.001>

Comentario a «Estudio de intervención para comprobar el efecto de la música clásica en directo durante hemodiálisis sobre la calidad de vida de pacientes con enfermedad renal crónica»

Comments: «Intervention study to verify the effect of live classic music during hemodialysis on the quality of life of patients with chronic kidney disease»

Sr. Director,

Hemos leído con mucho interés el artículo de Serrano et al.¹, en el cual ponen de manifiesto que la música en directo,

durante el tratamiento de hemodiálisis, tiene efectos positivos en la vida de los pacientes con enfermedad renal crónica. El ensayo se centra en la evaluación de variables descriptivas

