

4. Alonso Gomez AM. ¿Qué debe de conocer el nefrólogo de la afectación cardiaca del paciente en diálisis peritoneal? *Nefrología*. 2008;6:105-11.
5. Alhaj E, Alhaj N, Rahman I, Niaz T, Berkowitz R, Klapholz M. Uremic cardiomyopathy: an underdiagnosed disease. *Congest Heart Fail*. 2013;19:E40-5.
6. Pecoits-Filho R, Berberato SH. Echocardiography in chronic kidney disease: diagnostic and prognostic implications. *Nephron Clin Pract*. 2010;114:242-7.
7. Twardowski ZJ, Nolph KD, Prowant LB, Moore HL, Nielsen MP. Peritoneal equilibrium test. *Perit Dial Bull*. 1987;7:138-47.
8. Cueto-Manzano AM. Rapid solute transport in the peritoneum: physiologic and clinical consequences. *Perit Dial Int*. 2009;29:s90-5.
9. Chávez Valencia V, Orizaga de la Cruz C, Pazarin Villaseñor HL, Fuentes Ramírez F, Parra Michel R, Aragaki Y, et al. Frequency of peritoneal transport in a population of the Hospital General Regional No. 46, Instituto Mexicano del Seguro Social. *Gac Med Mex*. 2014;150:186-93.
10. Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr*. 2015;28:1-39.

Leonardo Pazarín-Villaseñor^{a,*}, Ulises Reyes-Lopez^b, Ana María de León-Flores^c, Alejandra Guillermina Miranda-Díaz^d, Jorge Andrade-Sierra^a

^a Servicio de Nefrología, Hospital Civil de Guadalajara «Dr. Juan I. Menchaca», Guadalajara, Jalisco, Mexico

^b Servicio de Medicina Interna, Hospital Civil de Guadalajara «Dr. Juan I. Menchaca», Guadalajara, Jalisco, Mexico

^c Servicio de Cardiología; Hospital Civil de Guadalajara «Dr. Juan I. Menchaca», Guadalajara, Jalisco, Mexico

^d Departamento de Fisiología, Centro Universitario de Ciencias de la Salud, Universidad de Guadalajara, Guadalajara, Jalisco, Mexico

* Corresponding author.

E-mail address: leopazarin@hotmail.com

(L. Pazarín-Villaseñor).

2013-2514/© 2016 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.nefro.2016.11.016>

Lengthening time to hemostasis in hemodialysis. Factor XI deficiency (hemophilia C): A case report[☆]

Alargamiento del tiempo de hemostasia en hemodiálisis. Déficit del factor XI (hemofilia C). A propósito de un caso

Dear Editor,

Factor XI is a dimeric serine protease which gene is located in chromosome 4 and it is present in platelets and plasma. Factor XI is involved in the intrinsic coagulation pathway by activating factor IX.^{1,2} It is produced by the liver and circulates as homodimer in an inactive form.^{3,4} The literature describes few cases with a deficit of factor XI (hemophilia C) on hemodialysis (HD).³

We are presenting a patient, 77 years with CKD secondary to nephroangiosclerosis on HD since July 2012. He had hypertension, ischemic heart disease, peripheral vascular disease at the femoral popliteal level, atherothrombotic vasculocerebral event with residual right hemiplegia. He has a humero-cephalic AVF in the left arm since October 2015 that presented repeated problems of puncture, bleeding complications, mainly hematomas and an increased time for hemostasis. Additional tests revealed abnormal coagulation when the patient was not taking anticoagulation of anti

platelets aggregation medications: the INR was 1.54 and the TTPa > 180s that were partially reversed with vitamin K, but did not prevent the bleeding.

Tests to evaluate hemostasis defect revealed an alteration of the intrinsic activation pathway of blood coagulation (rAPTT 1,32), likely related to a moderate deficit of factor XI (56% of the activity). El rest of factors involved in the intrinsic pathway were within the normal range (factor VIII 224%, factor IX 121%; factor XIII 84%). Lupus anticoagulant, anticardiolipin antibodies and anti b2GP1 (IgG and IgM) were negative. Hematologists advised the administration of nasal desmopresin (10 mcg/fosa nasal), 30 min before the end of the HD session, which succeeded in achieving hemostasia, without anticoagulation of the circuit and no thrombotic event has occurred so far to date.

Factor IX inhibits the fibrinolysis through the inhibition thrombin dependent fibrinolysis.^{5,6} Factor IX deficiency (hemophilia C) is an uncommon autosomal inherited disease found more frequently in some populations such as Ashkenazi Jews. Clinical symptoms may be detected in both homozygotes

[☆] Please cite this article as: Valga F, Monzón T, Henriquez Palop F, Antón G. Alargamiento del tiempo de hemostasia en hemodiálisis. Déficit del factor XI (hemofilia C). A propósito de un caso. *Nefrología*. 2017;37:544-545.

and heterozygotes. Symptomatology includes bleeding episodes that use to be less severe than in Hemophilia A or B and therefore it does not require prophylaxis.³ In these patients, bleeding may occur depending on the levels of other coagulation factors such as factor VIII and the von Willebrand factor.¹ There is little information on the clinical impact of Factor IX deficiency in HD patients. One case from Japan,³ had a APTT_a >120 s at the time of the AVF placement but this patient, as opposed to our case, had bleeding at the site of needle puncture.

Desmopresin (1-deamina-9-darginina-vasopresina) is a synthetic analog of vasopressin that increases the release of factor VIII and the von Willebrand factor¹ from the endothelium improving platelet aggregation. Desmopresin is used for the treatment of Type 1 von Willebrand disease and hemophilia A⁷ and is given to prevent bleeding associated to renal biopsies.⁸ Some authors hypothesize that desmopresin could stimulate the release of stored factor XI,⁹ this would be in addition to the effect on von Willebrand and VIII factors (normal in this case). We believe that in patients with a mild deficit is difficult to anticipate the severity of bleeding events since there is only a weak correlation with serum levels of factor XI.^{1,3,6,9} Other therapeutic options are transfusion of fresh frozen plasma, administration of factor XI concentrated after viral inactivation or the use of antifibrinolytic medication and the desmopresin combined with other therapies.^{1,6,9,10}

In summary, this case illustrates the need for investigating the cause of bleeding disorders in HD patients without assuming that it could be too much anticoagulation or to the addition of antiagregant therapy. This will improve the patient health since bleeding episodes will be avoided and will make the HD procedure safer.

REFERENCES

1. Bolton P. Trastornos de la coagulación poco comunes. In: Schulman S, editor. Tratamiento de la hemofilia. 39. Montreal, Quebec, Canada: World Federation of Hemophilia; 2006.
2. Butenas S, Dee JD, Mann KG. The function of factor XI in tissue factor-initiated thrombin generation. *J Thromb Haemost.* 2003;1:2103-11.
3. Takamizawa Y, Araki M, Yoshida N, Yoshioka T, Miura K. A case of a severe factor XI deficiency in patient undergoing hemodialysis without the use of heparin. *Blood Coagul Fibrinolysis.* 2014;25:898-9.
4. Wu W, Sinha D, Shikov S, Yip CK, Walz T, Billings PC, et al. Factor XI homodimer structure is essential for normal proteolytic activation by factor XIIa, thrombin, and factor XIa. *J Biol Chem.* 2008;283:18655-64.
5. Lowenberg EC, Meijers JC, Monia BP, Levi M. Coagulation factor XI as a novel target for antithrombotic treatment. *J Thromb Haemost.* 2010;8:2349-57.
6. Schumacher WA, Luetgen JM, Quan ML, Seiffert DA. Inhibition of factor XIa as a new approach to anticoagulation. *Arterioscler Thromb Vasc Biol.* 2010;30:388-92.
7. Siew DA, Mangel J, Laudenbach L, Schembri S, Minuk L. Desmopressin responsiveness at a capped dose of 15 mug in type 1 von Willebrand disease and mild hemophilia A. *Blood Coagul Fibrinolysis.* 2014;25:820-3.
8. Manno C, Bonifati C, Torres DD, Campobasso N, Schena FP. Desmopressin acetate in percutaneous ultrasound-guided kidney biopsy: a randomized controlled trial. *Am J Kidney Dis.* 2011;57:850-5.
9. Franchini M, Manzato F, Salvagno GL, Montagnana M, Lippi G. The use of desmopressin in congenital factor XI deficiency: a systematic review. *Ann Hematol.* 2009;88:931-5.
10. Duga S, Salomon O. Congenital factor XI deficiency: an update. *Semin Thromb Hemost.* 2013;39:621-31.

Francisco Valga*, Tania Monzón, Fernando Henriquez Palop, Gloria Antón

Centro de Diálisis Avericum Dr. Negrín, Las Palmas, Spain

* Corresponding author.

E-mail address: fvalga@hotmail.com (F. Valga).

2013-2514/© 2016 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). <https://doi.org/10.1016/j.nefro.2017.04.017>