

Consensus document

Consensus document for ultrasound training in the speciality of Nephrology[☆]

Maite Rivera Gorrín^a, R. Haridian Sosa Barrios^a, Carlos Ruiz-Zorrilla López^b, Juan Manuel Fernández^c, Silvia Marrero Robayna^c, José Ibeas López^d, Mercedes Salgueira Lazo^e, María Jesús Moyano Franco^e, Carlos Narváez Mejía^f, Manuel Ceballos Guerrero^f, Jordi Calabia Martínez^g, Antonio Luis García Herrera^h, Ramón Roca Teyⁱ, Vicente Paraíso Cuevas^j, José Luis Merino Rivas^j, Iyad Abuward Abu-Sharkh^k, Angels Betriu Bars^l, en representación del Grupo de Trabajo en Nefrología Diagnóstica e Intervencionista (GNDI) de la Sociedad Española de Nefrología (SEN)

^a Hospital Ramón y Cajal, Madrid, Spain

^b Hospital Río Carrión, Palencia, Spain

^c Hospital Dr. Negrín, Las Palmas de Gran Canaria, Spain

^d Parc Taulí Hospital Universitari, Institut d'Investigació i Innovació Parc Taulí I3PT, Universitat Autònoma de Barcelona, Sabadell, Barcelona, Spain

^e Hospital Macarena, Sevilla, Spain

^f Hospital Puerta del Mar, Cádiz, Spain

^g Hospital Josep Trueta, Girona, Spain

^h Hospital Puerto Real, Cádiz, Spain

ⁱ Hospital de Mollet, Mollet del Vallés, Barcelona, Spain

^j Hospital del Henares, Coslada, Madrid, Spain

^k Hospital Clínico Universitario, Santiago de Compostela, La Coruña, Spain

^l Institut de Recerca Biomèdica de Lleida (IRBLleida), Lleida, Spain

ARTICLE INFO

Article history:

Received 16 October 2019

Accepted 10 May 2020

Keywords:

Ultrasound

Vascular access

ABSTRACT

Ultrasound is an essential tool in the management of the nephrological patient allowing the diagnosis, monitoring and performance of kidney intervention. However, the usefulness of ultrasound in the hands of the nephrologist is not limited exclusively to the ultrasound study of the kidney. By ultrasound, the nephrologist can also optimize the management of arteriovenous fistula for hemodialysis, measure cardiovascular risk (mean intima thickness), implant central catheters for ultrasound-guided HD, as well as the patient's volemia using basic cardiac ultrasound, ultrasound of the Cava Inferior vein and lungs.

DOI of original article:

<https://doi.org/10.1016/j.nefro.2020.05.008>.

[☆] Please cite this article as: Gorrín MR, Barrios RHS, López CR-Z, Fernández JM, Robayna SM, López JI et al. Documento de Consenso para la Formación en Ecografía en la especialidad de Nefrología. *Nefrología*. 2020;40:623–633.

2013-2514/© 2020 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

From the Working Group on Interventional Nephrology of the Spanish Society of Nephrology (G.N.D.I. of the S.E.N.) we have prepared this Consensus Document that summarizes the main applications of ultrasound to Nephrology, including the necessary basic technical requirements, the framework normative and the level of training of nephrologists in this area. The objective of this work is to promote the inclusion of ultrasound, both diagnostic and interventional, in the usual clinical practice of the nephrologist and in the Nephrology Services portfolio with the final objective of offering diligent, efficient and comprehensive management to the nephrological patient.

© 2020 Sociedad Española de Nefrología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Documento de Consenso para la Formación en Ecografía en la especialidad de Nefrología

R E S U M E N

Diagnostic and interventional nephrology
Central line insertion
Renal replacement therapy
Renal biopsy

La ecografía es una herramienta esencial en el manejo del paciente nefrológico permitiendo el diagnóstico, seguimiento y realización de intervencionismo sobre el riñón. La utilidad de los ultrasonidos en Nefrología no se circunscribe exclusivamente al estudio ecográfico del riñón. Mediante ecografía el nefrólogo puede, además, optimizar el manejo de la fístula arteriovenosa para hemodiálisis, medir el riesgo cardiovascular (grosor íntima media), implantar catéteres centrales para hemodiálisis ecoguiados y ayudar en la colocación de los peritoneales, así como calcular la volemia del paciente mediante ecografía cardiaca básica, ecografía de la vena cava inferior y pulmonar.

Desde el G.N.D.I. de la S.E.N hemos elaborado este documento de consenso en el que se resumen las principales aplicaciones de la ecografía en Nefrología, incluyendo los requisitos técnicos básicos necesarios, el marco normativo y el nivel de capacitación de los nefrólogos en esta materia. El objetivo de este trabajo es promover la inclusión de la ecografía, tanto diagnóstica como intervencionista, en la práctica clínica habitual del nefrólogo y en la cartera de servicios de Nefrología con el objetivo final de ofrecer un manejo diligente, eficiente e integral al paciente nefrológico.

© 2020 Sociedad Española de Nefrología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Diagnostic and interventional nephrology (DIN), developed since its initiation by nephrologists. This is the area of Nephrology that includes all the imaging and interventional techniques used by the nephrologist: insertion of catheters as vascular access for renal replacement therapy (temporary and permanent catheters for haemodialysis [HD], placement of peritoneal dialysis catheters, renal biopsy, renal (kidneys, ureters, bladder) ultrasound and arteriovenous fistula [AVF]). DIN arose out of the necessity to solve the needs and difficulties encountered during routine clinical practice, in both diagnosis and treatment. Since ultrasound was first introduced into medicine in the 1950s,¹ its use has spread to multiple fields, including Nephrology, where it now plays a central role. By the end of the twentieth century, various authors had already written about the great utility of ultrasound in the management of renal patients.^{2,3}

Compared to other areas of Nephrology, DIN has remained relegated to the background from the point of view of patient care and teaching. Until recently, there was no recognised

specific training in the different techniques, and there is no official evaluation test or certification to assess the proficiency of physicians in performing these techniques.^{4,5}

There are a number of publications showing how the creation of a DIN section in the Nephrology department is highly efficient, improves the quality of care, reduces waiting times, improves patient safety, optimises resources, and is economically viable, preventing the overload of other departments.^{6–11} It also has an effect on the survival and viability of patients' vascular access for renal replacement therapy and on diligence in doctor and patient decision-making.

The Sociedad Española de Nefrología (SEN) [Spanish Society of Nephrology] therefore should promote and guarantee access to specific training in ultrasound for all nephrologists interested in performing these techniques.^{4,6,7} The SEN needs to support the development of learning tools¹² and establish routine practice standards. This should be accompanied by assessment and/or certification to officially validate the training undertaken by nephrologists and/or nephrology departments for performing diagnostic and interventional procedures. The SEN should also recommend the introduction

and systematic use of this technique for any of its indications to all nephrology departments.

Methods

On 10 October 2016, at the SEN Congress, a panel of experts, the Grupo de Nefrología Diagnóstica e Intervencionista (GNDI) [Diagnostic and Interventional Nephrology Group], met in Oviedo with the aim of encouraging and creating a document setting out the foundations of DIN for further development, teaching and quality control here in Spain. This panel of experts included nephrologists with a long track record and consolidated experience in DIN in their routine clinical practice.

To prepare this position paper, the experts drew up a discussion outline and questions based on the available data (systematic review) and the most important clinical evidence, debating each fundamental point until arriving at a consensus.

Each expert was responsible for writing an assigned section. The sections were put together and edited by MRG and RHSB in a draft, which was then approved by all panel members and presented as a consensus document. The conclusions are set out below.

Equipment and physical space required

Nephrology departments should have an ultrasound machine capable of performing studies in two-dimensional and Doppler modes. The equipment should have at least two probes or transducers: a low-frequency convex one for the exploration of the abdominal cavity (kidneys, bladder, prostate, liver-bile duct and abdominal aorta); and a high-frequency linear one for exploration of more superficial structures (pleura-lung, parathyroid glands, carotid and femoral arteries), and exploration and cannulation of the jugular and femoral veins. The same linear transducer will also be used in puncture and exploration of the vascular access for HD.

Certain applications require specific software; for example, for the measurement of carotid intima-media thickness (IMT), plus M mode and a low-frequency sector probe for echocardiography (Fig. 1).

Ultrasound scans can be performed in any physical space, even at the bedside, as the equipment is easily transportable. Low/minimal lighting is required to optimally display the image. Ultrasound-guided interventionism, in addition to a dimly lit room, requires a height-adjustable stretcher, ideally with wheels to evacuate the patient in the event of complications, a good light source, surgical and pharmacological material according to the procedure being carried out, and nursing staff support.

The intervention suite should have computer equipment to record the activity and compile the essential clinical reports of each examination (ultrasound and interventionism).

The Nephrology department must ensure storage and custody of the images obtained in a database, so they can be reviewed and, if necessary, compared with succes-

sive examinations of the same patient, and also to allow cases to be discussed with other healthcare professionals and transmission of knowledge. If the hospital uses electronic medical records, the images should be stored with the reports in the corresponding computer storage system.

Ultrasound applied to nephrology

A. Diagnostic ultrasound

Renal, bladder and prostate ultrasound

B-mode ultrasound of the kidney, bladder and prostate is a basic tool in the assessment of both acute and chronic kidney injury. It is essential in the study of renal function deterioration in the absence of previous glomerular filtration data. It enables obstructive causes and, using colour Doppler, vascular causes to be confirmed or ruled out quickly and non-invasively for the patient.^{2,13} The nephrologist needs to be able to assess³ the presence and location of both kidneys, their size, symmetry and echo structure, dilation of the urinary tract, and the presence of masses, cysts or lithiasis. For the bladder, we need to be able to recognise the presence of exophytic lesions, diverticula and imprints on the bladder wall, as well as assess the wall's thickness. Inside the bladder, we have to be able to recognise the presence of catheters or debris, as well as its filling volume. A full bladder and post-voiding examination should be performed to assess the presence of significant residue.

In the prostate, we have to be able to assess prostate size by measuring its three diameters in order to calculate the volume (Fig. 2).

The Doppler mode allows us to study intra- and extra-renal vessels, determine velocities for the diagnosis and follow-up of renal artery stenosis, and study vascular malformations, arterial or venous thrombosis and post-renal biopsy vascular complications.

Using colour Doppler (qualitative) we can assess the arterial and venous perfusion of the entire parenchyma, studying the renal artery and vein in all its sections. With spectral Doppler (quantitative) we can record the spectral wave of the vessel studied. From this record, we can make measurements of various parameters of vascular flow. In the indirect study, recordings are made of the intra-renal vessels, their morphology is analysed (e.g. to identify a "parvus et tardus" waveform, suggestive of stenosis), and the resistive, pulsatility and acceleration indices are calculated. The direct study (on the renal artery) is used to detect the presence of stenosis, measuring the velocities directly in the renal artery and in the aorta. The direct study is more complicated than the indirect one, as it is very difficult to visualise the renal artery in its entirety. In both the direct and indirect methods, a correct angle of insonation (angle between the ultrasound beam and the direction of blood flow) of a maximum of 60° is required for velocity measurements to be reliable (Fig. 3). Lastly, although in kidney transplantation the use of Doppler is mandatory, as there is a vascular anastomosis, in the native kidney there are only very specific indications (Fig. 4).

Essential

- Ultrasound with M, B and Doppler modes
 - High-frequency linear transducer
 - Low-frequency convex transducer
- Technology suite

Optional

- Contrast-enhanced ultrasound software
 - Software for measuring IMT
 - Sector transducer for echocardiography
 - Possibility of performing invasive techniques in the operating theatre
- C-arm X-ray machine

DIN: diagnostic and interventional nephrology IMT: intima-media thickness

Fig. 1 – Equipment required for DIN.**Renal ultrasound**

- Measurement of kidney size (longitudinal, transverse and anteroposterior)
- Measurement of kidney volume (L+T+AP/0.49)
- Parenchymal thickness
- Renal symmetry
- Kidney location, renal ectopia
- Rule out single kidney
- Study of urinary tract

Discover additional images and describe them: cysts, stones, kidney masses, etc.

Bladder ultrasound

- Measure bladder volume at maximum filling and post-voiding (residual volume)
 - Rule out wall abnormalities: diverticula, exophytic masses, etc.
 - Rule out abnormalities in interior: lithiasis, detritus, etc.
- Observe ureteric jet (colour Doppler)

Prostate ultrasound

- Measurement of prostate size (longitudinal, transverse and anteroposterior)
- Measurement of prostate volume (L+T+AP/0.49)

Fig. 2 – Data to be collected in a renal, bladder and prostate ultrasound.

- Studies renal vascularisation
 - Direct method: measures renal flow velocities in the main renal artery
 - Indirect method: measures velocities and resistive index in renal parenchyma
 - Types: colour Doppler (qualitative) and spectral Doppler (quantitative: spectral log analysis)
 - The study angle must be less than 60°
- Patient fasting: to avoid abdominal gas interference

Fig. 3 – Renal Doppler ultrasound: key aspects.**Ultrasound of vascular access. From pre-surgical mapping to diagnosis of complications**

Good vascular access (VA) is essential for the success of HD treatment, as it determines the quality of the dialysis, and complications can lead to high morbidity and mortality rates and increased healthcare costs.¹⁴

Preoperative mapping of the arterial and venous territory in the arms of patients who are candidates for VA, especially in patients with comorbidity, increases the rate of native VA and improves survival. Although arm mapping should be used in all patients, it is particularly necessary in cases where the habitual physical examination may be insufficient due to obesity, absent pulses, previous surgery, possible arterial disease

(advanced age, diabetes, cardiovascular disease) or possible venous disease (multiple previous venous cannulation).

The examination includes grey-scale B-mode ultrasound followed by a Doppler scan. Both arterial and venous exploration is from the distal area, starting at the wrist, to the proximal area, ending at the subclavian and axillary veins.¹⁴

The clinical management of VA can be optimised by the nephrologist's use of Doppler, as it enables early diagnosis of VA problems that may go unnoticed, helping to determine the degree of therapeutic urgency¹⁵ and avoiding unnecessary aggressive tests. It is therefore essential that the nephrologist knows the theoretical principles and the practical application of VA ultrasound.

- A.- Suspected renal vascular hypertension
- 1.- Acute onset or rapid progression of hypertension at any age, beginning before age 50 and severe hypertension in paediatric patients and young adults.
 - 2.- Confirmed hypertension refractory to properly managed medical treatment.
 - 3.- Deterioration in renal function induced by ACE inhibitor.
 - 4.- Hypertension associated with deterioration in renal function with no apparent cause.
 - 5.- Significant asymmetry (>1.5 cm) of kidney size in imaging tests.
- B.- Suspected ischaemic nephropathy
CKD of unknown cause in a patient with large vessel vascular disease

Fig. 4 – Renal Doppler ultrasound: indications in native kidney.

The nephrologist should perform serial Doppler scans of the VA (morphological surveillance and haemodynamic study)^{16,17} in order to detect significant stenosis,¹⁸ monitor non-significant stenosis,¹⁹ assess AVF maturation¹⁸ and diagnose various disorders, such as thrombosis,²⁰ aneurysms and pseudoaneurysms,^{18,20} and personalise the treatment of steal syndrome²¹ or its cardiac impact.^{22–24}

Arterial ultrasound (carotid, femoral and abdominal aorta): assessment of vascular risk in kidney disease

Cardiovascular (CV) disease is the leading cause of death in patients with chronic kidney disease (CKD), and the main culprit is atheromatosis. It is a chronic, progressive inflammatory process of the arteries, causing a thickening of the arterial intima-media layer until the formation of atheroma plaques (thickness >1.5 mm),²⁵ and is accelerated by traditional CV risk factors (age, sex, hypertension, diabetes, dyslipidaemia, smoking, etc.) and non-traditional factors (in patients with CKD: uraemic toxins, oxidative stress, chronic inflammation, anaemia, etc.).²⁶ Arterial ultrasound allows the direct and precise assessment of the damage caused by risk factors on the vascular wall, it being more accurate as it is not based solely on traditional CV risk factors and scores which are not validated in the population with CKD.²⁷ For that reason it is recommended in several different clinical practice guidelines.^{28,29} The diagnosis of subclinical atheromatous disease will help to detect the population most at risk and allow us to act early to prevent progression and CV events.

Ultrasound also makes it possible to assess the burden of atheromatous disease (number of territories and plaque area), analyse the composition of the plaques (lipids, fibrous tissue, calcium) and assess progression. As it is a systemic disease, it is important to explore all the accessible arterial territories: carotid (common, bifurcation or bulb, internal and external), femoral (common and superficial) and aorta (at the level of the abdomen). Various studies have shown that the prevalence of atheromatosis in femoral arteries, in the absence of carotid involvement, is in the range of 17%–20%.³⁰

Pleuropulmonary ultrasound

Lung ultrasound is a non-invasive way to detect lung congestion. It is of great help in the management of renal patients with dyspnoea, even in early subclinical stages. It is very useful in the assessment and monitoring of the response to treatment, in the prevention of volume overload in hypovolaemic heart patients treated with fluid therapy, and in the diagno-

sis and follow-up of pneumonia.^{31,32} The diagnosis of volume overload is made by evaluating ultrasound artefacts with diagnostic value such as lung rockets (multiple B lines) or comet tail artefacts.^{33,34}

Ultrasound is the best technique for the detection of pleural effusion; it is far superior to conventional radiology, as it is capable of detecting effusions of as little as 5 ml, compared to 150 ml in the posterior/anterior chest X-ray. The ultrasound pattern allows us to distinguish transudative from exudative. Lastly, ultrasound enables rapid detection of pneumothorax without the need to radiate the patient and with high reproducibility.

Transthoracic echocardiogram. Measurement of hydration status: ultrasound of the inferior vena cava

Transthoracic echocardiography is recognised for its utility in renal patients³⁵ and nephrologists should know how to perform it at a basic level.

Essentially, they need to be able to recognise pericardial effusion and left ventricular hypertrophy, and to know how to calculate the ejection fraction. A low-frequency sector probe is required. The degree of inspiratory collapse of the inferior vena cava measured by ultrasound can help estimate patients' effective volume status and, in conjunction with the symptoms, be a support in decision-making for the management of fluid therapy.³⁶

Parathyroid ultrasound

Ultrasound was the first imaging technique used to study the parathyroid glands. This is an easy technique for the nephrologist, with a quick and useful learning curve in the routine study of patients with secondary hyperparathyroidism, which provides great benefits and profitability.³⁷ Using ultrasound, we can determine the number and location of the parathyroid hypertrophy in order to plan a parathyroidectomy. The assessment of the parathyroid mass (volume) is of important prognostic value, as any increase is associated with a poorer response to drug treatment.³⁸ In some situations, ultrasound has been used to perform a chemical parathyroidectomy by intraglandular injection of alcohol/calcitriol.³⁹

Ultrasound in peritoneal dialysis. Exit site and tunnel, epigastric artery detection and catheter implantation

Ultrasound is highly useful in the implantation of the peritoneal catheter by the nephrologist, as it allows the systematic examination of the subcutaneous tissue, musculature and

fascia of the rectus muscles, and the location of the epigastric artery to avoid the risk of laceration and of the parietal peritoneum and the peritoneal cavity. It also makes it possible to locate the intestinal loops to ensure they are not in the catheter's path and to rule out adhesions between them.^{40–42} Lastly, an ultrasound scan to assess the bladder before implantation of the peritoneal catheter can help to prevent perforation in patients with post-void residual volume, with these patients being catheterised, if necessary, to ensure complete bladder emptying. Abdominal ultrasound also helps us guide the insertion of the catheter between the layers of the peritoneum,⁴³ observe how the infusion of peritoneal fluid separates the loops, and facilitate direction of the guide towards the pelvis,⁴² enabling correct placement of the catheter. It is important to use the ultrasound to identify the tip of the catheter in the pouch of Douglas before surgical closure at the end of the procedure.

In the post-implantation period, ultrasound in the hands of experienced personnel is very useful for assessing tunnel problems and complements other imaging techniques, such as plain X-ray or tomography, and can even sometimes replace them.⁴⁴ Ultrasound is useful for checking for infection or peri-catheter abscesses,^{44,45} bleeding or haematoma post-mechanical traction or post-implantation of the catheter, extrusion of the peritoneal catheter, peri-catheter and peritoneal cavity leaks⁴⁶ or catheter entrapment by omentum.⁴⁷

Finally, ultrasound can also be used to determine the thickness of the peritoneal membrane and how it relates to the dialysis technique's effectiveness, possibly avoiding the need for future aggressive tests, such as peritoneal biopsy.⁴⁸

Ultrasound of gastric motility

Gastroparesis or slowed gastric emptying is a known complication of diseases such as diabetes or amyloidosis, and also postoperatively, such as after kidney transplantation. Gastric emptying time is reportedly lengthened in patients undergoing any of the three types of renal replacement therapy compared to healthy subjects.⁴⁹ Gastroparesis can cause anorexia and malnutrition. It can be diagnosed by two-dimensional real-time ultrasound, taking dynamic images of gastric peristaltic activity and its effect on stomach emptying, and then calculating the total postprandial gastric volume.⁵⁰

Hepatobiliary ultrasound

Ultrasound allows early detection of fatty liver, a highly prevalent disorder in the general population and also, therefore, in renal patients. The echogenicity of the healthy renal parenchyma is very similar to that of the healthy liver. In fatty liver, the liver is whiter (hyperechogenic) than the renal parenchyma.⁵¹ Although previously considered a benign condition, progression to liver fibrosis and even cirrhosis has been reported,⁵² owing to which these patients should be referred to Gastroenterology for assessment.

With ultrasound, we can easily detect gallbladder sludge and gallstones, whether asymptomatic or complicated with cholecystitis, if we know how to recognise their ultrasound signs.

B. Ultrasound-guided nephrological intervention

Renal biopsy

Percutaneous renal biopsy (PRB) is one of the main procedures in the diagnosis of many different disorders, both of native and transplanted kidneys, and is currently the most widely used interventional technique in Nephrology departments.⁵³ With PRB not being free of complications,⁵⁴ the use of ultrasound when performing the procedure led to a clear reduction in post-biopsy complications, ranging from 10% to 15% of all patients biopsied using other procedures, to 4% to 6% after the introduction of ultrasound-guided PRB.⁵⁵

We can differentiate between three uses of ultrasound in renal biopsy:

- Dummy run to assess structure, characteristics of the kidney, access difficulties and possible contraindications to performing the biopsy (e.g. single kidney).
- Performing the PRB per se, with ultrasound enabling the exact puncture point to be chosen, the visualisation of the needle inside the kidney throughout the procedure, and the structures to avoid such as colon, large vessels or renal cysts.
- And afterwards, to detect possible complications, such as bleeding, haematoma, arteriovenous or arteriocholecal fistulas and pseudoaneurysms,^{56,57} even at the patient's bedside.

When the nephrologist is able to perform the PRB, this affords them autonomy and reduces the hospital stay by not having to depend on other departments. Acquiring skills in ultrasound-guided PRB requires specific training, experience and mastery in the morphological analysis of the healthy kidney and its pathological variables, and in the use of the Doppler mode for the analysis of post-biopsy vascular complications.

Implantation of tunnelled and non-tunnelled catheters for haemodialysis

Implantation of a catheter in a central vein with subcutaneous tunnelling is a valid alternative in various clinical situations requiring VA for HD, such as contraindication for an AVF, patients awaiting an AVF, or patients with an AVF, but not sufficiently developed to be used.

For the implantation of a catheter in a central vein, whether tunnelled or not, the AV Guidelines recommend ultrasound-guided puncture rather than implantation by anatomical reference 18, as ultrasound identifies the position of each vessel and its anatomical relationship, and reduces the risk of unwanted puncture of other adjacent vascular structures or the pneumothorax. It also enables detection of thrombosis, which would rule out the vessel for catheter implantation. In short, the use of ultrasound makes catheterisation easier, quicker and safer, and reduces both the likelihood of complications^{58,59} and the costs.⁶⁰ Ultrasound-guided puncture is ideal for cannulation of the jugular vein. However, for cannulation of the subclavian vein, the ultrasound view is limited by the interposition of the clavicle. For the implantation of a tunnelled catheter into a central, internal jugular or subclavian vein, it is recommended that after the puncture and the introduction of the guidewire the position of

Table 1 – Technical competencies of the interventional nephrologist in each of the domains that define this subspecialty of Nephrology.

Domain	Function	Competencies
Vascular ultrasound	Explore the patient's vascular tree using B-mode and Doppler ultrasound	<ul style="list-style-type: none"> • Explore the peripheral vascular tree, perform vascular access monitoring and diagnose vascular access complications • Use of ultrasound for targeted puncture for vascular access • Use of ultrasound to explore central venous vessels, diagnose complications in these vessels, and interpret the patient's hydration status
Renal ultrasound	Perform diagnostic ultrasound on native kidneys	<ul style="list-style-type: none"> • Diagnose the presence of disease in native kidneys and excretory system using B-mode ultrasound • Diagnose the presence of disease in renal vessels of native kidneys using Doppler ultrasound
	Perform diagnostic ultrasound on transplanted kidneys	<ul style="list-style-type: none"> • Diagnose the presence of disease in transplanted kidneys and excretory system using B-mode ultrasound • Diagnose the presence of disease in renal vessels of transplanted kidneys using Doppler ultrasound
Basic non-renal, non-vascular ultrasound	Perform basic ultrasound scan	<ul style="list-style-type: none"> • Detect and assess the presence of disease by B-mode and Doppler ultrasound in abdominal vessels, liver and bile duct • Assess hydration status. Perform pleuropulmonary ultrasound scan • Assess size of parathyroid glands. Perform parathyroid ultrasound scan
Echocardiography	Perform basic echocardiogram	<ul style="list-style-type: none"> • Detect and assess the presence of cardiac abnormalities and pericardial effusion using B-mode ultrasound
Renal biopsy	Perform real-time ultrasound-guided renal biopsy	<ul style="list-style-type: none"> • Apply the ultrasound-guided renal biopsy technique to native and transplanted kidneys, guaranteeing the safety of the technique. • Diagnose complications derived from the technique using B-mode and Doppler ultrasound
Catheters for haemodialysis	Perform ultrasound-guided cannulation of central vessels and implant catheters for haemodialysis	<ul style="list-style-type: none"> • Perform ultrasound-guided implantation of temporary haemodialysis catheters in central vessels, guaranteeing the safety of the technique • Perform ultrasound-guided implantation of tunnelled haemodialysis catheters in central vessels, guaranteeing the safety of the technique
	Implant tunnelled catheters in central vessels under scope control	<ul style="list-style-type: none"> • Use the scope to cannulate central vessels and implant tunnelled haemodialysis catheters, guaranteeing the safety of the technique • Have a full understanding of the necessary radioprotection measures for both the patient and healthcare professionals
	Remove permanent tunnelled catheters	<ul style="list-style-type: none"> • Use a semi-surgical technique to remove and replace tunnelled catheters, guaranteeing the safety of the technique
Catheters for peritoneal dialysis	Implant peritoneal catheters	<ul style="list-style-type: none"> • Implant peritoneal catheters ensuring the safety of the technique by using ultrasound for assessment and positioning • Use ultrasound to diagnose peritoneal catheter-related complications
	Remove peritoneal catheters	<ul style="list-style-type: none"> • Use a semi-surgical technique to remove and replace peritoneal catheters, guaranteeing the safety of the technique

Table 2 – Consolidated courses in Diagnostic and Interventional Nephrology in Spain.

Name	Year started	Director	Site
[0,1–4] <i>Ultrasound and Interventionism</i> Masters' in Diagnostic and Interventional Nephrology	2011	M.E. Rivera Gorrín	Hospital Universitario Ramón y Cajal [Ramón y Cajal University Hospital] (Madrid) Universidad de Alcalá de Henares [Alcalá de Henares University]
Seminar on applied techniques in haemodialysis by Nephrology	2014	V. Paraíso, J.L. Merino	Hospital Universitario del Henares [El Henares University Hospital]
Course on ultrasound-guided interventional techniques in Nephrology	2017	D. del Pino y Pino	IAVANTE - Fundación Progreso y Salud, CMAT (Complejo Multifuncional Avanzado de Simulación e Innovación Tecnológica [Advanced Multifunctional Complex for Simulation and Technological Innovation]), Granada
Theoretical-practical course on interventionism in renal patients	2017	J. Ibeas	Parc Taulí Hospital Universitari [Parc Taulí University Hospital] (Sabadell)
[0,1–4] [0,1–4] <i>Ultrasound in vascular access</i> Course on vascular access for haemodialysis	2009	R. Álvarez Lipe, M.T. González Álvarez, A. Foraster Roselló	Hospital Universitario Gonzalo Blesa [Gonzalo Blesa University Hospital] (Zaragoza)
Theoretical-practical course of ultrasound for vascular access in haemodialysis	2009	J. Ibeas	Parc Taulí Hospital Universitari [Parc Taulí University Hospital] (Sabadell)
Hands-on Skills Training Workshop in Dialysis Access: A Multidisciplinary Approach	2011	J. Ibeas	Parc Taulí Hospital Universitari [Parc Taulí University Hospital] (Sabadell)
Parc Taulí International Vascular Access Symposium	2014	J. Ibeas	Parc Taulí Hospital Universitari [Parc Taulí University Hospital] (Sabadell)
Theoretical and practical course: multidisciplinary approach to vascular access for haemodialysis	2016	A Vilar Gimeno	Hospital General Universitario de Valencia [Valencia University General Hospital]
[0,1–4] [0,1–4] <i>Ultrasound in Nephrology</i> Theoretical-practical course on ultrasound applied to Nephrology	2016	M.E. Rivera Gorrín	Hospital Universitario Ramón y Cajal [Ramón y Cajal University Hospital] (Madrid)
Theoretical-practical course on renal ultrasound and Doppler	2017	J. Calabia	Hospital Universitari Doctor Josep Trueta [Doctor Josep Trueta University Hospital] (Girona)
5th theoretical-practical course on ultrasound for nephrologists: clinical ultrasound applied to problems in Nephrology	2017	C. Narváez Mejía	Sociedad Andaluza de Nefrología [Andalusian Society of Nephrology]

the guidewire be checked by radioscopia to confirm that the intravascular end is lodged in the right place, and to avoid catheter dysfunction owing to malpositioning. 18 Radioscopia also facilitates the identification of thrombosis or stenosis in central veins with the use of contrast, helping to rule out the affected vessel. Radioscopic confirmation is not considered necessary if the catheter is implanted in the femoral vein.

Implantation of peritoneal catheter

As already discussed in the *Ultrasound in peritoneal dialysis* section, ultrasound will be very useful in peritoneal catheter implantation in order to optimise placement, avoid

complications such as puncture of the epigastric artery or intestinal loops, and to confirm the exact position.

Ultrasound-guided puncture of the arteriovenous fistula

Puncture of the AVF can be difficult due to poor maturation, small diameter, juxta-anastomotic stenosis or, around the puncture site, proximity to arteries or nerves, existence of collaterals, previous unsuccessful punctures with haematomas, or even not knowing the direction of cannulation in the case of deep AVF. The characteristics of the HD programme population, with a progressive increase in age and vascular comorbidity, have led to an increase in this type of puncture with extra difficulties.

To counteract these difficulties, Doppler ultrasound performed at the patient's bedside by the nephrologist and/or haemodialysis unit nurse can increase the sensitivity of early detection of abnormalities, resulting in reduced morbidity and mortality rates.^{61–63} Doppler provides information on the patency, flow, depth and diameter of the vessel when performing the targeted puncture,^{64,65} and allows the path of the needle to be tracked within the vessel.^{66–70} This last aspect is essential, as it allows the needle to be repositioned during the HD session if necessary.

Although there are limited studies on the utility of Doppler ultrasound for ultrasound-guided puncture of the VA, its recent incorporation into routine clinical practice has shown that it can reduce the number of punctures, facilitate cannulation in difficult-to-access AVF, and reduce patient discomfort deriving from the puncture. It is therefore necessary to include strategies aimed at training professionals in this specific area.⁷¹

Competencies of the interventional nephrologist

The technical competencies that the interventional nephrologist needs to acquire in each of the domains that define this subspecialty of Nephrology are listed in [Table 1](#).

The following are the *basic or mandatory* areas that the nephrologist must have a mastery of:

- Renal and bladder/prostate ultrasound, both 2-D and Doppler modes.
- AVF ultrasound and ultrasound-guided puncture of the same.
- Cannulation of temporary HD catheters, both jugular and femoral.
- Ultrasound-guided renal biopsy, both native and transplanted kidneys.

The following would be *secondary* or *optional* areas:

- Cardiopulmonary ultrasound.
- Vascular ultrasound: carotid, femoral and abdominal aorta.
- Vascular mapping of the upper limb prior to AVF.
- Gastrointestinal ultrasound: gastric and hepatobiliary.
- Parathyroid ultrasound.
- Implantation of tunnelled catheters for HD.
- Implantation of catheters for peritoneal dialysis{2}

Regulatory and administrative framework

The training programme for the speciality of Nephrology (ORDER SCO/2604/2008. BOE [Official State Gazette] 15/09/2008) in point 5.3.4 establishes renal ultrasound, renal biopsy, temporary vascular access and peritoneal catheters as within the knowledge, skills and development of investigation, diagnosis and treatment techniques.⁷² In the new training programme, currently under development, Interventional Nephrology is included among the skills that the nephrologist must master. Lastly, in the 2016–2020 SEN strate-

gic plan, the Governing Board included reassessment of the speciality of Nephrology as a priority, defending its own competencies and developing emerging areas such as Interventional Nephrology.⁷³

Last of all, we, as members of the GNDI, believe it is necessary for the SEN to establish an accreditation and training system in ultrasound applied to Nephrology in order to guarantee quality and homogeneous training and clinical practice in this area for Spanish nephrologists. While there are no accredited centres, Spanish nephrologists have a wide range of courses available to them, the majority accredited and with SEN endorsement, in addition to university degrees. [Table 2](#) shows a list of the most consolidated courses in Spain.

Conflicts of interest

The authors declare that they have no conflicts of interest.

REFERENCES

1. Ludwig, G.D. and Struthers, F.W. Considerations underlying the use of Ultrasound to detect Gallstones and Foreign Bodies in Tissue. Naval Medical Research Institute Reports, Project #004 001, Report No. 4, June 1949.
2. Rivera M. Incorporación de la ecografía a la práctica rutinaria del nefrólogo: nuestra experiencia. *Nefrología*. 1995;15:104–7.
3. O'Neill WC. Renal ultrasound: A procedure for nephrologists. *Am J Kidney Dis*. 1997;30:579–85.
4. Sachdeva M, Ross DW, Shah HH. Renal ultrasound, catheter placement, and kidney biopsy experience of US Nephrology fellows. *Am J Kidney Dis*. 2016;68(2):187–92.
5. Berns JS, O'Neill WC. Performance of procedures by nephrologists and Nephrology fellows at US Nephrology training programs. *Clin J Am Soc Nephrol*. 2008;3:941–7.
6. Asif A, Byers P, Vieira CF, Roth D. Developing a comprehensive diagnostic and interventional nephrology program at an academic center. *Am J Kidney Dis*. 2003;42:229–33.
7. Rasmussen RL. Establishing an interventional nephrology suite. *Semin Nephrol*. 2002;22:237–41.
8. Beathard GA, Litchfield T. Effectiveness and safety of dialysis vascular access procedures performed by interventional nephrologists. *Kidney Int*. 2004;66:1622–32.
9. Asif A, Merrill D, Briones P, Roth D, Beathard GA. Hemodialysis vascular access: percutaneous interventions by nephrologists. *Semin Dial*. 2004;17(6):528–34.
10. Gadallah MF, Pervez A, El-Shahawy MA, et al. Peritoneoscopic versus surgical placement of peritoneal dialysis catheters: A prospective randomized study on outcome. *Am J Kidney Dis*. 1999;33:118–22.
11. Asif A. Peritoneal dialysis access-related procedures by nephrologists. *Semin Dial*. 2004;17(5):398–406.
12. Gorrin Rivera, Correa Gorospe C, Burguera V, Ortiz Chercoles AI, Liaño F, Quereda C. Teaching innovations in ultrasound guided renal biopsy. *Nefrología*. 2016;36:1–4.
13. Meola M, Petrucci I. Ultrasound and color Doppler in Nephrology. *Acute Kidney Injury*. *G Ital Nefrol*. 2012;29:599–615.
14. Roca-Tey R, Arcos E, Comas J, Cao H, Tort J, Catalan Renal Registry C. Starting hemodialysis with catheter and mortality risk: persistent association in a competing risk analysis. *The journal of vascular access*. 2016;17(1):20–8.

15. Ibeas J, Vallespin J. Ecografía del acceso vascular para hemodiálisis: conceptos teóricos y prácticos. *Criterios. Nefrología*. 2012;3(6):15.
16. Malik J, Slavikova M, Svobodova J, Tuka V. Regular ultrasonographic screening significantly prolongs patency of PTFE grafts. *Kidney international*. 2005;67(4):1554-8.
17. Scaffaro LA, Bettio JA, Cavazzola SA, Campos BT, Burmeister JE, Pereira RM, et al. Maintenance of hemodialysis arteriovenous fistulas by an interventional strategy: clinical and duplex ultrasonographic surveillance followed by transluminal angioplasty. *Journal of ultrasound in medicine*. 2009;28(9):1159-65.
18. Ibeas J, Roca-Tey R, Vallespin J, Moreno T, Monux G, Marti-Monros A, et al. Spanish Clinical Guidelines on Vascular Access for Haemodialysis. *Nefrología*. 2017;37 Suppl 1:1-191.
19. Tuka V, Slavikova M, Krupickova Z, Mokrejsova M, Chytilova E, Malik J. Short-term outcomes of borderline stenoses in vascular accesses with PTFE grafts. *Nephrology, dialysis, transplantation*. 2009;24(10):3193-7.
20. Wiese P, Nonnast-Daniel B. Colour Doppler ultrasound in dialysis access. *Nephrology, dialysis, transplantation*. 2004;19(8):1956-63.
21. Beathard GA, Spergel LM. Hand ischemia associated with dialysis vascular access: an individualized access flow-based approach to therapy. *Seminars in dialysis*. 2013;26(3):287-314.
22. Wijnen E, Keuter XH, Planken NR, van der Sande FM, Tordoir JH, Leunissen KM, et al. The relation between vascular access flow and different types of vascular access with systemic hemodynamics in hemodialysis patients. *Artificial organs*. 2005;29(12):960-4.
23. Agarwal AK. Systemic Effects of Hemodialysis Access. *Advances in chronic kidney disease*. 2015;22(6):459-65.
24. Vaes RH, Tordoir JH, Scheltinga MR. Systemic effects of a high-flow arteriovenous fistula for hemodialysis. *The journal of vascular access*. 2014;15(3):163-8.
25. Touboul PJ, Hennerici MG, Meairs S, Adams H, Amarenco P, Bornstein N, et al. Mannheim carotid intima-media thickness and plaque consensus (2004-2006-2011). An update on behalf of the advisory board of the 3rd, 4th and 5th watching the risk symposia, at the 13th, 15th and 20th European Stroke Conferences, Mannheim, Germany, 2004, Brussels, Belgium, 2006, and Hamburg, Germany, 2011. *Cerebrovasc Dis*. 2012;34:290-6.
26. Betriu-Bars A, Fernández-Giráldez E. La ecografía carotídea en el diagnóstico precoz de enfermedad arterial ateromatosa en la enfermedad renal crónica. *Nefrología*. 2012;32:7-11.
27. Coll B, Betriu A, Martínez-Alonso M, Borràs M, Craver L, Amoedo ML, et al. Cardiovascular risk factors underestimate atherosclerotic burden in chronic kidney disease: usefulness of non-invasive tests in cardiovascular assessment. *Nephrol Dial Transplant*. 2010;25:3017-25.
28. Piepoli MF, Hoes AW, Agewall S, Albus Ch, Brotons C, Catapano AL, et al. 2016 European Guidelines on cardiovascular disease prevention in clinical practice. *Cerebrovasc Dis*. 2012;34:290-6.
29. Mancia G, Fagard R, Narkiewicz K, Redón J, Zanchetti A, Bo H, et al. The Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *European Heart Journal*. 2016;37:2315-81.
30. Fernández-Friera L, Peñalvo JL, Fernández-Ortiz A, Ibañez B, López-Melgar B, Laclaustra M, et al. Prevalence, Vascular Distribution, and Multiterritorial Extent of Subclinical Atherosclerosis in a Middle-Aged Cohort. The PESA (Progression of Early Subclinical Atherosclerosis) Study. *Circulation*. 2015;131:2104-13.
31. Siriopol D, Hogas S, Voroneanu L, Onofriescu M, Apetrii M, Oleniuc M, et al. Predicting mortality in haemodialysis patients: a comparison between lung ultrasonography, bioimpedance data and echocardiography parameters. *Nephrol Dial Transplant*. 2013;28:2851-9.
32. Blaiwas M. Lung Ultrasound in Evaluation of Pneumonia. *J Ultrasound Med*. 2012;31:823-6.
33. de la Quintana Gordon FB, Alcorta N, Fajardo Pérez M. Ecografía pulmonar básica. Parte 2. Patología Parenquimatosa. *Rev Esp Anestesiología Reanim*. 2015;62:337-49.
34. Mallamaci F, Benedetto FA, Tripepi R, Rastelli S, Castellino P, Tripepi G, Picano E, Zoccali C. Detection of Pulmonary Congestion by Chest Ultrasound in Dialysis Patients. *JACC: Cardiovascular imaging*. 2010;3:586-94.
35. García Fernández MA, Zamorano JL, García Robles JA. Manual de Ecocardiografía: Indicaciones e interpretación en la práctica clínica. Editorial EDIMED; 2004, <http://secardiologia.es/images/stories/la-sec/latinoamerica/manual-ecocardiografia.pdf>.
36. Ciozda W, Kedan I, Kehl DW, Zimmer R, Khandwalla R, Kimchi A. The efficacy of sonographic measurement of inferior vena cava diameter as an estimate of central venous pressure. *Cardiovasc Ultrasound*. 2015;14:33.
37. Torregrosa JV, Félez I, Fuster D. Utilidad de las técnicas de imagen en el hiperparatiroidismo secundario. *Nefrología*. 2010;30:158-67.
38. Restrepo C. Ultrasound detection of parathyroid hyperplasia and correlation with clinical and laboratory findings in patients with chronic kidney disease. *Rev Colomb Radiolog*. 2011;22:3341-7.
39. Nakamura M, Fuchinoue S, Teraoka S. Clinical experience with percutaneous ethanol injection therapy in hemodialysis patients with renal hyperparathyroidism. *AmJ Kidney Dis*. 2003;42:739-45.
40. Maya ID. Ultrasound/fluoroscopy-assisted placement of peritoneal dialysis catheters. *Semin Dial*. 2007;20:611-5.
41. Díaz-Mancebo R, Del Peso-Gilsanz G, Bernabeu D, Bajo-Rubio MA, Selgas-Gutierrez R. Colocación de catéter peritoneal guiado por ecografía. *NefroPlus*. 2015;7:98.
42. Doñate T. Guías de diálisis peritoneal y la práctica diaria. *Nefrología* 25. Suplemento 2:33-38.
43. Balaskas EV, Ikononopoulos D, Sioulis A, et al. Survival and complications of 225 catheters used in continuous ambulatory peritoneal dialysis: one-center experience in Northern Greece. *Perit Dial Int J Int Soc Perit Dial*. 1999;19 Suppl 2:S167-71.
44. Díaz-Mancebo R, Del Peso-Gilsanz G, Bernabeu D, Bajo-Rubio MA, Selgas-Gutierrez R. Utilidad de la ecografía en el diagnóstico de los problemas mecánicos en pacientes en diálisis peritoneal. *Nefroplus*. 2015;7:99-101.
45. Espósito F, Di Serafino M, Ambrosio C, Rita Panico M, Malacario F, Mercogliano C, et al. Chronic peritoneal dialysis in children: the role of ultrasound in the diagnosis of peritoneal catheter obstruction. *J Ultrasound*. 2016;19:191-6.
46. Latich I, Luciano RL, Mian A. Image-Guided Approach to Peritoneal Dialysis Catheter Placement. *Tech Vasc Interv Radiol*. 2017;20:75-81.
47. Golay V, Trivedi M, Roychowdhary A, Arora P, Sarkar D, Singh A, et al. Ultrasound-guided CAPD catheter insertion. *Perit Dial Int*. 2013;33:454-8.
48. Guz G, Bali M, Poyraz NY, Bagdatoglu O, Yeğin ZA, Doğan I, et al. Gastric emptying in patients on renal replacement therapy. *Ren Fail*. 2004;26:619-24.
49. Darwiche G, Almér LO, Björgell O, Cederholm C, Nilsson P. Measurement of Gastric Emptying by Standardized Real-Time Ultrasonography in Healthy Subjects and Diabetic Patients. *J Ultrasound Med*. 1999;18:673-82.
50. Anwar Javed, Aamir Muhammad Omer, Sanaullah Zeeshanul Hasnain Imdad, Parveen Ishrat, Yousaf Nasreen. Can

- ultrasound abdomen help in early diagnosis of diabetes mellitus? An observational study. *J Ayub Med Coll Abbottabad*. 2015;27:807-10.
51. Musle Acosta M, Bolaños Vaillant S, Gómez García Y, Toirac Romani CA, Rodríguez Cheong M. Alteraciones ecográficas en pacientes con diabetes mellitus [artículo en línea]. *MEDISAN*. 2008;12(3), <http://bvs.sld.cu/revistas/san/vol12_3.08/san04308.htm> [consulta: 15.01.2018].
 52. Stephen A, Harrison MD. Liver Disease in Patients With Diabetes Mellitus. *J Clin Gastroenterol*. 2006;40:68-76.
 53. Registro de glomerulonefritis de la Sociedad Española de Nefrología.
 54. KDIGO Clinical Practice for glomerulonephritis. *Kidney International Supplements*. 2012;2:8-12.
 55. Hergesell O, Felten H, Andrassy K, et al. Safety of ultrasound-guided percutaneous renal biopsy, retrospective analysis of 1090 consecutive cases. *Nephrol Dial Transplant*. 1998;13:975-7.
 56. Whittier LW, Kobrabet SM. Timing of complications in percutaneous renal biopsy. *J Am Soc Nephrol*. 2004;15:142-7.
 57. Rivera Gorrín M. Biopsia renal ecodirigida. *Nefrología*. 2010;30:490-2.
 58. Aydin Z, Gursu M, Uzun S, Karadag S, Tatli E, Sumnu A, et al. Kazancioglu R Placement of hemodialysis catheters with a technical, functional, and anatomical viewpoint. *Int J Nephrol*. 2012;2012:302826.
 59. Rabindranath KS, Kumar E, Shail R, Vaux EC. Ultrasound use for the placement of haemodialysis catheters. *Cochrane Database Syst Rev*. 2011;11:CD005279.
 60. Bowen MC, Mone MC, Nelson EW, Scaife CL. Image-guided placement of long-term central venous catheters reduces complications and cost. *The Am J Surgery*. 2014;208:937-41.
 61. Hernández-López J. La enfermería de diálisis con la ecografía en los accesos vasculares para hemodiálisis. <http://www.sedyt.org/docs/cursos/comunicaciones-viii-curso-accesos-vasculares/ecografia-accesos-vasculares-dr-hernandez.pdf> (consulta: 5.02.2018).
 62. Patel RA, Stern AS, Brown M, Bhatti S. Bedside ultrasonography for arteriovenous fistula cannulation. *Semin Dial*. 2015;28:433-4.
 63. Ward F, Faratro R, McQuillan RF. Ultrasound-Guided Cannulation of the Hemodialysis Arteriovenous Access. *Semin Dial*. 2017;30:319-25.
 64. Kamata T, Tomita M, Iehara N. Ultrasound-guided cannulation of hemodialysis access. *Renal Replacement Therapy*. 2016;2(7).
 65. Ibeas López J, Vallespín-Aguado J. Ecografía del acceso vascular para hemodiálisis: conceptos teóricos y prácticos. *Criterios. Nefrología Sup Ext*. 2012;3:21-35.
 66. Egan G, Healy D, O'Neill H, Clarke-Moloney M, Grace PA, Walsh SR. Ultrasound guidance for difficult peripheral venous access: systematic review and meta-analysis. *Emerg Med J*. 2013;30:521-6.
 67. Hanafusa N, Noiri E, Nangaku M. Vascular access puncture under ultrasound guidance. *Ther Apher Dial*. 2014;18:213-4.
 68. Jiménez Almonacid P. Fístulas arteriovenosas para hemodiálisis. En: Lorenzo V, López Gómez JM (Eds) *Nefrología al Día*. <http://www.revistanefrologia.com/es-monografias-nefrologia-dia-articulo-fistulas-arteriovenosas-hemodialisis-38> (consulta 5.02.2018).
 69. Garu Pueyo C, Granados Navarrete I, Moya Mejía C, García Blanco M, Vinuesa García-Ciaño X, Ramírez Vaca J, et al. La punción del acceso vascular en hemodiálisis es una necesidad, el método Buttonhole una opción. *Rev Soc Esp Enferm Nefrol*. 2011;14:30-6.
 70. Hernández López J. Punción con ecografía dirigida de la fístula arteriovenosa dificultosa. *Dial Traspl*. 2011;32:126-7.
 71. Marticorena RM, Mills L, Sutherland K, McBride N, Kumar L, Bachynski JC, Rivers C, Petershofer EJ, Hunter J, Luscombe R, Donnelly S. Development of competencies for the use of bedside ultrasound for assessment and cannulation of hemodialysis vascular access. *CANNT J*. 2015;25:28-32.
 72. <http://www.senefro.org/modules.php?name=webstructure&idwebstructure=21>.
 73. http://www.senefro.org/contents/webstructure/Plan_estrategico_de_la_S.E.n.pdf.