

## Impact of the SEN-SEMFYC consensus document on chronic kidney disease upon organisation of our nephrology departments

*Nefrología* 2008; 28 (3) 350

**To the editor:** The challenge posed by a pathology such as chronic kidney disease (CKD)<sup>1-3</sup> imposes an adequate coordination of the nephrology departments and primary care (PC) centres.<sup>4</sup> The recently issued «SEN-SEMFYC consensus document on chronic kidney disease» represents a significant effort in this direction.

A concern arising when a standardised procedure to manage CKD is prepared is the potential saturation of nephrology outpatient clinics. In our experience, implementation of a common procedure with PC led to a 40% increase in the number of first visits in the first year. This has led us to be cautious about criteria to be agreed on. As regards values of the glomerular filtration rate estimated by the MDRD formula (eGFR), the document proposes referral of patients with

eGFR < 45 mL/min/1.73 m<sup>2</sup> in those older than 70 years and with eGFR < 30 mL/min/1.73 m<sup>2</sup> in patients older of 70 years. Are these the most reasonable criteria for all departments?

In order to analyse their impact on the Alcañiz health sector (Teruel), covering an aged rural population of 83,456 inhabitants, we used our database of 18,922 patients over 18 years of age who were performed laboratory tests at our healthcare centres during 2006. CKD prevalence was 16.4%, and 208 patients would meet these referral criteria. Use of values proposed in our protocol (< 65 years, eGFR < 45 mL/min/1.73 m<sup>2</sup>; 65-80 years, eGFR > 30 mL/min/1.73 m<sup>2</sup>; > 80 years, eGFR < 20 mL/min/1.73 m<sup>2</sup>) would decrease this figure to 113 patients. The characteristics of both groups are shown in the table.

Result analysis suggests that in our health sector we could use the referral criteria in the consensus document in patients under 70 years of age, but always stressing that morbidity and mortality are particularly increased in males under 60 years of age with eGFR ranging from 45-60 mL/min/1.73 m<sup>2</sup>, who should be considered at a high cardiovascular risk.<sup>2</sup> However, application of such criteria would not be feasible to older pa-

tients, particularly those over 80 years of age. The excessive patient volume and the limited prevention of cardiovascular and renal risk we could achieve lead us to reconsider the question.<sup>5</sup> In this case, adequate patient management and referral of patients with progressive kidney function impairment (since replacement therapy is often feasible) or requiring management of their complications (particularly anaemia) would be left at PC discretion.

Based on these data, we think that training and cooperation with PC and adjustment of criteria for referral to nephrology outpatient clinics based on the capacity of each department are essential for early detection and follow-up of CKD. Adequate management of CKD by PC in male patients younger than 60 years with eGFR values ranging from 45-60 mL/min/1.73 m<sup>2</sup> and in those older than 75-80 years with eGFRs ranging from 20-30 mL/min/1.73 m<sup>2</sup> is particularly relevant.

- Otero A, Gayoso P, García F, De Francisco AL, on behalf of the EPIRCE study group: epidemiology of chronic renal disease in the Galician population: results of the pilot Spanish EPIRCE study. *Kidney Int* 2005; (Supl. 99): S16-S19.
- Go AS, Chertow GM, Fan D, McCulloch CE, Hsu C-Y. Chronic kidney disease and the risk of death, cardiovascular events, and hospitalization. *New Engl J Med* 2004; 351: 1296-1305.
- O'Hare AM, Bertenthal D, Covinsky KE, Landefeld CS, Sen S, Mehta K, Steinman MA, Borzecki A, Walter LC. Mortality risk stratification in chronic kidney disease: One size for all ages? *J Am Soc Nephrol* 2006; 17: 846-854.
- Alcázar R, De Francisco AL. Acción estratégica de la SEN frente a la enfermedad renal. *Nefrología* 2006; 26: 1-4.
- Alcázar R, Egocheaga MI, Orte L, Lobos JM et al. Documento de consenso SEN-semFYC sobre la enfermedad renal crónica. *Nefrología* 2008; 28 (3): 273-281.
- Locatelli F, Pozzoni P. Chronic kidney disease in the elderly: is it really a premise for overwhelming renal failure? *Kid Int* 2006; 69: 2118-2120.

L. M. Lou Arnal, J. M.<sup>a</sup> Turón Alcaine, B. Campos Gutiérrez and B. Boned Juliani  
 Servicios de Nefrología. Dirección de Atención Primaria y Bioquímica del Hospital de Alcañiz. Unidad de Investigación del Sector Sanitario de Alcañiz. Grupo Aragonés de Investigación en Atención Primaria REDIAPP.

**Table I. Characteristics of the population groups that would require referral to nephrology outpatient clinics according to the different criteria for the glomerular filtration rate estimated by the MDR formula**

	Prediction according to the SEN-SEMFYC document	Prediction according to Alcañiz protocol
Study population	18,922	18,922
No. of patients meeting referral criteria	208 (1.1%)	113 (0.6%)
Sex (male/female)	37.5%/62.5%	42.5%/57.5%
Mean age	72.9 ± 14	67.1 ± 14
A. Age groups		
< 65 years	50 (24%)	50 (44.3%)
65-80 years	79 (38%)	47 (41.6%)
> 80 years	79 (38%)	16 (14.1%)
B. Age groups		
< 70 years	89 (42.8%)	56 (49.6%)
> 70 years	119 (75.2%)	57 (50.4%)
PCr (mg/dL)	2.19 ± 0.98	2.39 ± 1.2
GFR (mL/min/1.73 m <sup>2</sup> )	30 ± 9.3	29.7 ± 11
K/DOQI stage		
- 3	80 (38.5%)	46 (40.7%)
- 4	116 (55.8%)	55 (48.7%)
- 5	12 (5.7%)	12 (10.6%)