

A) BRIEF PAPERS ABOUT BASIC RESEARCH OR CLINICAL EXPERIENCES

The risks of digoxin in the elderly

Nefrologia 2010;30(1):131-2

Dear Editor,

The elderly represent 17% of the population and 70% of drug expenditure. Adverse drug events account for between 7.2 and 14% of hospital admissions for these patients in internal medicine in Spain.¹ Over the last decades this consumption has increased from 3 to 4-8.2 drugs/day,² and it has been shown that the quantity of drugs ingested is related to adverse effects. The prevalence of hidden kidney disease in old age, and internal changes in patients due to the pathology of multiple medication has led to an increase in iatrogenic events. The following case of digitalis intoxication is submitted as an example of this.

An 82-year old woman with a history of heart failure episodes, with a previous serum Cr reference of 1.2mg/dL, arrived at an emergency ward in a mentally confused state. Multiply medicated, with an unknown dose of digoxin, plus diuretics, anticoagulants, etc., and no access to medical attention in the previous 4 months. On examination there were signs of moderate dehydration, psychomotor agitation, with a blood pressure of 120/60mmHg and a heart rate of 34 beats/min. At the emergency unit, biochemistry values were: Hct 42%, Hb 11g/dl, urea 199, Cr 4.8, K 6.8, digoxinaemia 5.4, pH 7.29, HCO 18. Chest radiograph, abdominal ultrasound and CT head scans were unremarkable. ECG: nodal rhythm of 34 per minute. She was given emergency haemodialysis without ultrafiltration, controlling the hyperkalaemia without any changes in the ECG. She was not considered in need for admission to the intensive care unit, nor were any other measures taken, such as insertion of a pacemaker or administration of digoxin antibody. For 48 hours the cardiovascular and neurological status of the patient remained unchanged, with subsequent resolution of the whole process after rehydration, digoxin levels

Table 1. Compliance of risk factors of digitalis toxicity in kidney function before and after treatment

	Yes	No	Possible
Advanced age	+		
Female sex	+		
Kidney disease			+
Use of diuretics	+		
Heart failure	+		
Digoxin dose			+

normalised, with recovery of urine output and mental state.

This case had several risk factors that increased the risk of drug and digoxin intoxication (Table 1), with the drug being responsible for the symptoms.

In order to evaluate kidney function when administering this drug in the elderly, more factors than the serum creatinine values should be taken into account.³ It is likely that this patient had hidden kidney disease. Our obligation as nephrologists is to carry out emergency dialysis, due to the presence of oliguric acute kidney injury after a pump failure with hyperkalaemia, obviously, not to decrease digoxin levels. The final outcome was satisfactory without any more aggressive measures being taken due to the history and age of the patient.

Digitalis toxicity is present in 0.4% of hospital admissions, with a toxicity of 10-18% in old peoples' homes, leading to a 34% morbidity and mortality of 1%. This toxicity may be chronic, sometimes accidental and also due to attempted suicide.⁴ Situations (e.g. kidney disease) or drugs that increase their levels (verapamil and amiodarone, among others) can lead to intoxication. Moreover, as is known, hyperkalaemia, acidosis and hypercalcaemia enhance the effect.

Apart from the known clinical condition, hyperkalaemia requires a

special mention as a prognostic factor and indicator of treatment. This may result in not only decreased excretion due to kidney disease, but a dysfunction of the Na-K-ATPase due to the digitalis effect. This would be a first order prognostic marker because it faithfully reflects the magnitude of the deleterious effect, and it is an indication for treatment with digoxin antibody (K > 5mEq/dL). Intravenous calcium administration to correct the electrocardiographic effect mediated by hyperkalaemia must be avoided, due to the risk of further increasing intracellular calcium linked to digitalis toxicity.

In another sense, arrhythmia therapy indications, among others, should be addressed by the relevant specialty, and we must practically confine ourselves to the repeated administration of atropine, if necessary, as there is a parasympathetic hyperactivity that can influence the placement of pacemakers, performing gastric lavage, etc. The use of digoxin antibodies seems far more efficient and faster, although in kidney disease a rebound effect may occur after initial administration.⁵

1. Alcalde P, Dapena MD, Nieto MD, Fontecha BJ. Ingreso hospitalario atribuible a efectos adversos a medicamentos. Rev Esp Geriatr Gerontol 2001;36:340-4.
2. Blasco F, Martínez López de Letona J, Pérez Maestu R, Villares P, Ponce J. Estudio piloto sobre el consumo defármacos en ancianos que ingresan en un hospital. An Med Intern 2004;21:69-71.
3. Cepeda Piorno J, Pobes Martínez de

Salinas A, González García ME, Fernández Rodríguez E. Utilidad de la ecuación MDRD para detectar insuficiencia renal oculta y disminuir el riesgo de sobredosificación digitalica. *Nefrología* 2009;29(2):150-5.

4. Antman EM, Wenger TL, Butler VP, et al. Treatment of 150 cases of life-threatening digitalis intoxication with digoxin-specific Fab antibody fragments. Final report of a multicenter study. *Circulation* 1990;81:1744.
5. Mehta RN, Mehta NJ, Gulati A. Late rebound digoxin toxicity after digoxin-specific antibody Fab fragments therapy in anuric patient. *J Emerg Med* 2002;22:203.

A. Suárez Laurés, A. Pobes Martínez, L. Quiñones Ortiz, R. Forascepi

Nephrology Department.

Cabueñes Hospital, Gijón, Spain.

Correspondence: Luis Quiñones Ortiz
Servicio de Nefrología.

Hospital Cabueñes. Gijón. Spain.

luysquio@hotmail.com

Role of healthcare coordinator: experience in a haemodialysis satellite unit

Nefrología 2010;30(1):132

Dear Editor,

Chronic kidney disease (CKD) is a serious public health problem with biological, mental and social implications.^{1,4} The characteristics of CKD patients on haemodialysis (HD) have changed in recent years. Age and comorbidity have increased, which has implications on functional aspects. Haemodialysis centres are seeing patients with a significant degree of dependence.⁵ The greater percentage of elderly patients makes it more difficult to modify the dialysis technique, requiring increased social care.⁶ For these reasons, and to improve the quality of care given to our patients, the idea of creating the role of unit healthcare coordinator was suggested in 2006. Initially, the coordinator saw 5 patients on a part time basis, although this help proved so invaluable that he was recently

made full time. He is currently involved in managing tasks for a total of approximately 75 patients. Among his duties are liaison with the referral hospital, processing additional test documentation and consultations with other specialist departments. He also accompanies patients, if necessary, thus preventing the continuous loss of appointments and helps the nephrologist communicate with other specialists. Sometimes he acts as liaison with primary care and, if necessary, with social workers where there is a need for care. He also organises transfers, holidays, etc., which is very helpful for patients who are involved in complex bureaucratic procedures, thus facilitating their adaptation within the limitations of their disease.

This results in patients in our unit benefitting from improved care that goes beyond haemodialysis sessions, with a more global view of the problems they experience.

We, as nephrologists, have seen a clear improvement in organisation and care, providing the quality health care required by our patients.

We therefore encourage other haemodialysis centres to consider appointing a similar healthcare coordinator among their personnel

1. Kimmel PL. Psychosocial factors in dialysis patients. *Kidney Int* 2001;59(5):1599-613.
2. Schieppati A, Remuzzi G. Chronic renal diseases as a public health problem: epidemiology, social, and economic implications. *Kidney Int* 2005;68(Supl. 98):S7-S10.
3. Eknayan G, Lameire N, Barsoum R, Eckardt KU, Levin A, Levin N, et al. The burden of kidney disease: improving global outcomes. *Kidney Int* 2004;66(4):1310-4.
4. Barsoum RS. Chronic kidney disease in the developing world. *N Engl J Med* 2006;354(10):997-9.
5. Arenas MD, Álvarez-Ude F, Angoso M, Berdud I, Antolín A, Lacueva J, et al. Valoración del grado de dependencia funcional de los pacientes en hemodiálisis

(HD): estudio multicéntrico. *Nefrología* 2006;26(5):600-8.

6. Martín de Francisco AL. Hemodiálisis en el anciano. *Nefrología* 1998;18(Supl. 4):10-4.

C. Rodríguez Adanero¹, S. Estupiñán Torres¹, R. Pérez Morales¹, S. García Rebollo², V. Lorenzo Sellarés²

¹Haemodialysis Satellite Unit, University Hospital of the Canary Islands, Spain.

²Nephrology Department, University Hospital of the Canary Islands, Spain.

Correspondence:

Concepción Rodríguez Adanero

Unidad Satélite de Hemodiálisis.

Hospital Universitario de Canarias. Spain.

cradanero@senefro.org

luysquio@hotmail.com

Ten years' experience of an Intensive and Continuous Theoretical/Practical Training Course in Peritoneal Dialysis

Nefrología 2010;30(1):132-3

Dear Editor,

Although peritoneal dialysis (PD) is now a recognised kidney replacement technique, it is still used much less than haemodialysis. According to SEN (Spanish Society of Nephrology) statistics (from the SEN), of 4,543 incident cases in replacement therapy during 2007, 12.4% were initiated with PD; of 36,388 prevalent cases, 46% started with haemodialysis (HD), 47.9% underwent transplantation and 6.1% received DP,¹ proportions similar to some European countries.²

Among the reasons for this, one very important is the lack of knowledge and expertise of many professionals, leading some to not offer PD as a possible dialysis option.

According to the General Health Law (*Law 14/1986, 25 April*), "each patient has the right to free choice between different treatment options". The